

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K X		A2	(11) International Publication Number: WO 92/18089 (43) International Publication Date: 29 October 1992 (29.10.92)
(21) International Application Number: PCT/US92/02237 (22) International Filing Date: 27 March 1992 (27.03.92) (30) Priority data: 682,809 9 April 1991 (09.04.91) US (60) Parent Application or Grant (63) Related by Continuation US 682,809 (CIP) Filed on 9 April 1991 (09.04.91) (71) Applicant (for all designated States except US): THE UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).			(72) Inventors; and (75) Inventors/Applicants (for US only) : ABRAHAM, Irene [US/US]; 1805 Waite, Kalamazoo, MI 49008 (US). HESTER, Jackson, Boling [US/US]; 9219 East ML Avenue, Galesburg, MI 49053 (US). (74) Agent: STEIN, Bruce; Corporate Patents & Trademarks, The Upjohn Company, Kalamazoo, MI 49001 (US). (81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, PL, RO, RU, SD, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US. Published Without international search report and to be republished upon receipt of that report.

(54) Title: USE OF AMINES TO SENSITIZE MULTIDRUG-RESISTANT CELLS



(57) Abstract

The sensitizing amines of the present invention, illustrated by the steroidal amines (I), the alkyl amines (II), bicyclic amines (III), bicyclic ethers (IV), tricyclic compounds (V), indoles (VI) and various species are useful in treating individuals who have cancer has become resistant to cancer chemotherapeutic agents and in preventing the resistance from developing or slowing the rate of resistance to the chemotherapeutic agents.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Tchad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

USE OF AMINES TO SENSITIZE MULTIDRUG-RESISTANT CELLS

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention is a method of sensitizing desensitized cancer cells to various
5 chemotherapeutic agents so as to better treat the cancer.

2. Description of the Related Art

It is known that cancer cells become resistant to some chemotherapeutic agents and are even resistant to many different chemotherapeutic agents. This is a significant impediment to cancer chemotherapeutic treatment.

10 It would be of great benefit to those who have cancers which can be treated with chemotherapeutic agents to have an agent which will sensitize the resistant cancer cells to the chemotherapeutic agents being used to treat the cancer.

Several classes of compounds which sensitize cancer cells to chemotherapeutic agents are known. One group of compounds are calcium channel blockers or related analogues, some
15 of which no longer act as calcium channel blockers. For example, some synthetic isoprenoids *Cancer Research* 46, 4453 (1986) and pyridine analogs *Int. J. Cancer* 45, 508 (1990), cause multidrug resistance reversal. Reserpine *Biochem. Pharmacol.* 30, 2191 (1981), quinidine and cyclosporine and related analogs *J. Clin. Invest.* 77, 1405 (1986) some without the original bioactivities, also sensitize multidrug resistance cells. Phenothiazines such as the calmodulin
20 inhibitors thioridazine, trifluoperazine and chlorpromazine *J. Nat. Cancer Inst.* 76, 839 (1986) and *Cancer Let.* 30, 251 (1986) sensitize multidrug resistant cells. Analogs of Vinca alkaloids and anthracyclines such as N-acetyldaunorubicin *Cancer Res.* 40, 1077 (1980) will also sensitize multidrug resistant cells, as will steroids *J. Biol. Chem.* 264, 782 (1989), and *Biochem. Biophys. Res. Commun.* 158, 1066 (1989)]. Other agents that sensitize multidrug
25 resistant cells include the antibiotic cefoperazone *Cancer Research* 49, 6901 (1989), tamoxifen *Cancer Res.* 44, 4392 (1984), vitamin A, *Br. J. Cancer* 56, 267 (1987) and chloroquine *Cancer Lett.* 30, 251 (1986).


At least two features seem to be of major importance in determining whether a compound can act as a sensitizer. One is lipophilicity *Cancer Res.* 50, 3997 (1990) and the other is
30 ability to modulate binding on the P-glycoprotein, *Advances In Pharmacol.*, 21, 185 (1990). Several of the previously identified sensitizing compounds appear to act by competing for binding to the P-glycoprotein. The mode of action of others that do not compete for binding is unknown.

It has been found that various steroidal and non-steroidal amines also sensitize
35 desensitized cancer cells to the common chemotherapeutic agents. These sensitizing compounds

are useful in anti-cancer therapy to sensitize cancer cells to be killed by traditional cytotoxic drugs (chemotherapeutic agent). More particularly, 21-[4-2,6-di-1-pyrrolidinyl]-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione (tirilazad mesylate; 74,006) sensitizes desensitized cancer cells as well as or better than verapamil, which is a standard sensitizer. Verapamil has the disadvantage of causing cardiac toxicity in humans at doses that are required for *in vivo* sensitization of cancer cells.

SUMMARY OF INVENTION

Disclosed is a method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing steroidal amine of formula (I)



where:

20 (A-I) R_6 is α - R_{6-1} : β - R_{6-2} , R_{10} is α - R_{10-1} : β - R_{10-2} and R_7 is α -H: β -H, where one of R_{6-1} and R_{6-2} is -H, and the other is -H, -F, or C_1 - C_3 alkyl, R_{10-2} is -CH₃, R_{10-1} and R_5 taken together are $-(CH_2)_2-C(=R_{3-3})-CH=$ or $-CH=CH-CO-CH=$, where R_{3-3} is =O or α -H: β -OR₃₋₄ or α -OR₃₋₄: β -H, where R_{3-4} is -H, -CO-CH₃, -CO-C₂H₅, -CO-C₆H₅, -CO-O-CH₃ or -CO-O-C₂H₅;

25 (A-II) R_5 is α - R_{5-3} : β - R_{5-4} , R_6 is α - R_{6-3} : β - R_{6-4} , R_{10} is α - R_{10-3} : β - R_{10-4} and R_7 is α -H: β -H, where one of R_{6-3} and R_{6-4} is -H, and the other taken together with one of R_{5-3} and R_{5-4} forms a second bond between C₅ and C₆, R_{10-4} is -CH₃, R_{10-3} and the other of R_{5-3} and R_{5-4} taken together is $-(CH_2)_2-C(H)(OH)-CH_2-$;

30 (A-III) R_{10} and R_5 taken together are $=CH-CH=C(OR_3)-CH=$ where R_3 is -H, C_1 - C_3 alkyl, -CO-H, C_2 - C_4 alkanoyl or benzyl, R_6 is α - R_{6-5} : β - R_{6-6} where one of R_{6-5} and R_{6-6} is -H, and the other is -H, -F, or C_1 - C_3 alkyl and R_7 is α -H: β -H;

(A-IV) R_5 is α - R_{5-7} : β - R_{5-8} , R_6 is α - R_{6-7} : β - R_{6-8} , R_7 is α -H: β -H and R_{10} is α - R_{10-7} : β - R_{10-8} , where one of R_{5-7} and R_{5-8} is -H, R_{10-7} and the other of R_{5-7} and R_{5-8} taken together are $-(CH_2)_2-C(=R_{3-3})-CH_2$, where R_{3-3} is as defined above, R_{10-8} is -CH₃, where one of R_{6-7} and R_{6-8} is -H and the other is -H, -F, or C_1 - C_3 alkyl;

35 (A-V) R_6 is R_{6-9} : R_{6-10} , R_7 is R_{7-9} : R_{7-10} , R_{10} is α - R_{10-9} : R_{10-10} where one of R_{6-9}

-3-

and R_{6-10} is -H and the other taken together with one of R_{7-9} and R_{7-10} forms a second bond between C_6 and C_7 , and the other of R_{7-9} and R_{7-10} is -H, R_{10-10} is $-CH_3$, R_{10-9} and R_5 taken together are $-(CH_2)_2-C(=R_{3-3})-CH=$ or $-CH=CH-CO-CH=$, where R_{3-3} is as defined above;

5 where:

(C-I) R_{11} is $\alpha-R_{11-1}:\beta-R_{11-2}$, where one of R_{11-1} and R_{11-2} is taken together with R_9 to form a second bond between C_9 and C_{11} and the other of R_{11-1} and R_{11-2} is -H;

(C-II) R_9 is -Cl and R_{11} is =O or $\alpha-H:\beta-R_{11-4}$ where R_{11-4} is -Cl or -OH;

(C-III) R_9 is -H or -F and R_{11} is =O or $\alpha-R_{11-5}:\beta-R_{11-6}$, where one of R_{11-5} and R_{11-6} is -H, and the other of R_{11-5} and R_{11-6} is -H, -OH or C_1-C_{12} alkoxy;

(C-IV) R_9 is -H or -F and R_{11} is $\alpha-O-CO-R_{11-7}:\beta-H$, where R_{11-7} is

(A) C_1-C_3 alkyl,

(B) C_1-C_{12} alkoxy,

(C) furanyl,

15 (D) $-NR_{122}R_{123}$, where one of R_{122} and R_{123} is -H, methyl or ethyl and the other is -H, C_1-C_4 alkyl or phenyl,

(E) $-X_3-Aryl$, where X_3 is -O- or a valence bond, where Aryl is phenyl optionally substituted with 1 thru 2 -Cl, -Br, C_1-C_3 alkoxy, -COOH, -NH₂, C_1-C_3 alkylamino, di(C_1-C_3)alkylamino, where the alkyl groups are the same or different, 1-
20 pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2-C_4 acylamino and -NH-CHO or with 1 -F or -CF₃;

where:

(D-I) R_{16} is $R_{16-1}:R_{16-2}$ and R_{17} is $R_{17-1}:R_{17-2}$, where one of R_{16-1} and R_{16-2} is -H or $-CH_3$ and the other taken together with one of R_{17-1} and R_{17-2} forms a second bond
25 between C_{16} and C_{17} , and the other of R_{17-1} and R_{17-2} is $-C(=Z)-(CH_2)_n-$
 $NR_{21-\alpha}R_{21-\beta}$, where Z is =O, =CH₂ or $R_{17-9}:-H$ where R_{17-9} is -H or $-CH_3$, where n is 0 thru 6, where

(A) $R_{21-\alpha}$ is

(1) $-(CH_2)_m-NR_{21-1}-Heteroaryl$, where m is 2, 3 or 4, where R_{21-1} is -
30 H or C_1-C_3 alkyl, where Heteroaryl is:

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is

(i) -F,

(ii) -Cl,

35 (iii) -Br,

(iv) C_1-C_5 alkyl,

-4-

(v) $-\text{CH}_2-\text{CH}=\text{CH}_2$.

(vi) -Aryl, where Aryl is as defined above,

(vii) $-\text{NR}_{21-3}\text{R}_{21-3}$ where the R_{21-3} 's are the same or different and are -H, $\text{C}_1\text{-C}_3$ alkyl or $-\text{CH}_2-\text{CH}=\text{CH}_2$,

5 (viii α) $^*\text{CH}_2-(\text{CH}_2)_q-\text{CH}_2-\text{N}^*$ where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,

(viii β) $^*\text{CH}_2-\text{CH}_2-(\text{CH}_2)_c-\text{G}-(\text{CH}_2)_d-\text{CH}_2-\text{CH}_2-\text{N}^*$

where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4), where G is -O-, -S-, -SO-, -SO₂- or -NHR₂₁₋₄, where R_{21-4} is -H, $\text{C}_1\text{-C}_3$ alkyl, or Aryl as defined above, where c and d are the same or different and are 0 thru 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,

(ix) 3-pyrrolin-1-yl, (F-5)

(x) pyrrol-1-yl optionally substituted with $\text{C}_1\text{-C}_3$ alkyl, (F-6)

15

(xi) piperidin-1-yl optionally substituted with 1 or 2 $\text{C}_1\text{-C}_3$ alkyl, (F-7)

 C_3 alkyl,

(xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)

(xiii) 1-hexamethyleneimino containing a 3- or 4- double bond or 3- and 5- double bonds, (F-9)

20

bond or 3- and 5- double bonds,

(xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by two $\text{C}_1\text{-C}_3$ alkyl being the same or different, (F-10)

(xv) -OH,

(xvi) $\text{C}_1\text{-C}_3$ alkoxy,

25

(xvii) $-\text{NR}_{21-7}(\text{CH}_2)_e-\text{Q}$ where Q is 2-pyridinyl where R_{21-7} is -H or $\text{C}_1\text{-C}_3$ alkyl and e is 0 thru 3,

(xviii) pyridin-2-, 3- or 4-yl,

(b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the 4- and/or 6- position with R_{21-2} is as defined above, (F-11)

30

(c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2- and/or 6- position with R_{21-2} is as defined above, (F-12)

(d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1 or 2 R_{21-2} as is defined above, (F-13)

(e) pyrazin-2-yl optionally substituted with 1 or 2 R_{21-2} as is defined above, (F-14)

35

defined above,

(f) imidazol-2-yl optionally substituted in the 1 position with

C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-15)

(g) 1,2,4-triazol-3-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)

(h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)

(i) benzo[b]thien-2-yl, (F-18)

(j) indol-2-yl, (F-19)

(k) benzo[b]thiazol-2-yl, (F-20)

(l) benzimidazol-2-yl, (F-21)

(m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl]piperazinyl, (F-22)

(n) 1,2,4-triazol-3-yl optionally substituted at the 5- and/or 6-position with R_{21-2} as is defined above, (F-23)

(2) (1-piperazinyl)-(C_2-C_4)alkyl optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)

(4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is

(a) $-O-CH_2CH_2-Y$, where Y is C_1-C_3 alkylamino, di(C_1-C_3)alkylamino where the alkyl groups are the same or different, C_3-C_6 alkyleneimino, optionally substituted with 1 or 2 C_1-C_3 alkyl,

(b) $-NR_{21-20}CH_2CH_2-Y$, where R_{21-20} is -H or C_1-C_3 alkyl and Y is as defined above,

(c) $-(CH_2)_g-N(R_{21-20})-Heteroaryl$, where g is 2, 3 or 4, and where R_{21-20} and Heteroaryl are as defined above,

(5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1-C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3-C_6 heterocyclic ring and where m is as defined above,

(6) $-(CHCH_3)_b-(CH_2)_f-R_{21-24}$, where b is 0 and f is 1 thru 3 or b is one and f is 0 thru 3, where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1-C_3 alkoxy, $-NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1-C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4-C_7 cyclicamino ring,

(7) $-(CH_2)_i-Heteroaryl$, where i is 1 thru 4 and Heteroaryl is as defined above,

(8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl
where Heteroaryl is as defined above, (F-25)

(9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by
-Heteroaryl where Heteroaryl is as defined above, and (F-26)

5 (B) $R_{21-\beta}$ is

(1) -H,

(2) C_1-C_3 alkyl,

(3) C_5-C_7 cycloalkyl,

(4) $-(CH_2)_m-NR_{21-1}$ -Heteroaryl, where m, R_{21-1} and Heteroaryl are as
10 defined above,

(5) (1-piperazinyl)-(C_2-C_4)alkyl optionally substituted in the 4- position
with -Aryl or -Heteroaryl as defined above, (F-24)

(6) $-(CH_2)_m-X_4$, where m and X_4 are as defined above,

(7) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where m, R_{21-22} and R_{21-23} are as
15 defined above,

(8) $-(CHCH_3)_b-(CH_2)_f-R_{21-24}$, where b, f and R_{21-24} are as defined
above,

(C) $R_{21-\alpha}$ and $R_{21-\beta}$ are taken together with the attached nitrogen atom to form
a heterocyclic ring selected from the group consisting of

20 (1) 2-(carboxy)-1-pyrrolidinyl optionally as the C_1-C_3 alkyl ester or as a
pharmaceutically acceptable salt, (F-27)

(2) 2-(carboxy)-1-piperidinyl optionally as the C_1-C_3 alkyl ester or as a
pharmaceutically acceptable salt, (F-28)

(3) 2-(carboxy)-1-hexamethyleneimino optionally as the C_1-C_3 alkyl
25 ester or as a pharmaceutically acceptable salt, (F-29)

(4) 2-(carboxy)-1-heptamethyleneimino optionally as the C_1-C_3 alkyl
ester or as a pharmaceutically acceptable salt, (F-30)

(5) 1-piperazinyl substituted in the 4- position with $R_{21-28}CO-(CH_2)_j$ -
where R_{21-28} is -Aryl, $-NR_{21-29}Aryl$ and 2-furanyl, where R_{21-29} is -H or C_1-C_3 alkyl, where
30 j is 0 thru 3 and Aryl is as defined above, (F-31)

(6) 1-piperazinyl substituted in the 4- position with Heteroaryl- $(CH_2)_j$ -,
where Heteroaryl and j are as defined above, (F-32)

(7) 1-piperazinyl substituted in the 4- position with
Aryl- $(CH_2)_j$ -, where Aryl and j are as defined above, (F-33)

35 (8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as
defined above, (F-34)

(9) 1-piperazinyl substituted in the 4- position with Heteroaryl-NR₂₁₋₂₉-CO-(CH₂)_i, where Heteroaryl, R₂₁₋₂₉ and i are as defined above; (F-35)

(D-II) R₁₆ is α -R₁₆₋₃: β -R₁₆₋₄ where one of R₁₆₋₃ and R₁₆₋₄ is -H and the other is -H, -F, -CH₃ or -OH, and R₁₇ is =CH-(CH₂)_p-NR_{21- α} R_{21- β} , where p is 1 or 2, where R_{21- α} and R_{21- β} are as defined above;

(D-III) R₁₆ is α -R₁₆₋₅: β -R₁₆₋₆ and R₁₇ is α -R₁₇₋₅: β -R₁₇₋₆, where R₁₆₋₅ is -H, -OH, -F or -CH₃ and R₁₆₋₆ is -H, -OH, -F, or -CH₃, with the proviso that at least one of R₁₆₋₅ and R₁₆₋₆ is -H, where R₁₇₋₅ is -H, -OH, -CH₃, -CH₂CH₃, C₂-C₇ alkanoyloxy or -O-CO-Aryl, where Aryl is as defined above, and where R₁₇₋₆ is -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , where Z, n, R_{21- α} and R_{21- β} are as defined above;

(D-IV) the 16,17-acetonide of a compound where R₁₆₋₅ is -OH, R₁₆₋₆ is -H, R₁₇₋₅ is -OH and R₁₇₋₆ is -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , where Z, n, R_{21- α} and R_{21- β} are as defined above;

with the following overall provisos that:

(I) one of R₁₆₋₁ or R₁₆₋₂ is taken together with one of R₁₇₋₁ or R₁₇₋₂ to form a second bond between C₁₆ and C₁₇, only when R₁₀ is α -R₁₀₋₁: β -R₁₀₋₂, α -R₁₀₋₃: β -R₁₀₋₄, α -R₁₀₋₇: β -R₁₀₋₈ or α -R₁₀₋₉: β -R₁₀₋₁₀;

(II) R₁₇ is =CH-(CH₂)_p-NR_{21- α} R_{21- β} , only when R₁₀ is α -R₁₀₋₁: β -R₁₀₋₂, α -R₁₀₋₃: β -R₁₀₋₄, α -R₁₀₋₇: β -R₁₀₋₈ or α -R₁₀₋₉: β -R₁₀₋₁₀;

(III) R₅ and R₁₀ taken together are =CH-CH=C(OR₃)-CH=, only when R₁₇ is α -R₁₇₋₅: β -R₁₇₋₆ or the 16,17-acetonide of a compound where R₁₆ is α -OH: β -H and R₁₇ is α -OH: β -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , and

(IV) R₅ is α -R₅₋₇: β -R₅₋₈, only when R₁₇ is α -R₁₇₋₅: β -R₁₇₋₆ or α -OH: β -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , or the 16,17-acetonide thereof; and pharmaceutically acceptable salts thereof.

Also disclosed is a method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing alkyl amine of formula (II)



where:

n₂ is 3-14;

X₂ is -H,

-OH,

-O-CO-(C₁-C₄ alkyl),

-O-CO-H,

-O-CO-O-(C₁-C₄ alkyl),

(C₁-C₄) alkoxycarbonyl,

-O-CO-Aryl where Aryl is -φ optionally substituted with 1 thru 3 of the following which may be the same or different:

-OH,

5

-OCH₃,

-F, -Cl, -Br, -CF₃,

-C₁-C₃ alkyl, and

-CO-R₅ where R₅ is

-OH,

10

-NH₂,

-NHR₆ where R₆ is

-φ,

C₁-C₃ alkyl and

-N(R₁₄)(R₁₅) where R₁₄ and R₁₅ are the same or

15 different and are C₁-C₃ alkyl,

-O-Aryl, where Aryl is as defined above,

-CH(OH)Aryl, where Aryl is as defined above,

Aryl, where aryl is as defined above;

(A) R_{21-α} is

20

(1) -(CH₂)_m-NR₂₁₋₁-Heteroaryl, where m is 2, 3 or 4, where R₂₁₋₁ is -H or C₁-C₃ alkyl, where Heteroaryl is:

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R₂₁₋₂, being the same or different, where R₂₁₋₂ is

(i) -F,

25

(ii) -Cl,

(iii) -Br,

(iv) C₁-C₅ alkyl,

(v) -CH₂-CH=CH₂,

(vi) -Aryl, where Aryl is phenyl optionally substituted with 1

30 through 2 -F, -Cl, -Br, C₁-C₃ alkoxy, -COOH, -NH₂, C₁-C₃ alkylamino, di(C₁-C₃)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C₂-C₄ acylamino and -NH-CHO or with 1 -F or -CF₃;

(vii) -NR₂₁₋₃R₂₁₋₃ where the R₂₁₋₃s are the same or different

35 and are -H, C₁-C₃ alkyl or -CH₂-CH=CH₂,

(viii) *CH₂-(CH₂)_q-CH₂-N* - where the atoms marked with an

asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 through 5,

- (viii) $^*\text{CH}_2-(\text{CH}_2)_c-\text{G}-(\text{CH}_2)_d-\text{CH}_2-\text{N}^*$ -where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4),
 5 where G is -O-, -S-, -SO-, -SO₂- or -NR₂₁₋₄-, where R₂₁₋₄ is -H, C₁-C₃ alkyl, or Aryl as defined above, where c and d are the same or different and are 0 through 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,
- (ix) 3-pyrrolin-1-yl, (F-5)
 (x) pyrrol-1-yl optionally substituted with C₁-C₃
 10 alkyl, (F-6)
 (xi) piperidin-1-yl optionally substituted with 1 or 2
 C₁-C₃ alkyl, (F-7)
 (xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)
 (xiii) 1-hexamethyleneimino containing a 3- or 4- double bond
 15 or 3- and 5- double bonds, (F-9)
 (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by
 two C₁-C₃ alkyl being the same or different, (F-10)
 (xv) -OH,
 (xvi) C₁-C₃ alkoxy,
 20 (xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where R₂₁₋₇
 is -H or C₁-C₃ alkyl and e is 0 through 3,
 (xviii) pyridin-3- or 4-yl,
 (xix) -CF₃,
 (xx) -CCl₃,
 25 (xxi) -SCH₃,
 (b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the
 4- and/or 6- position with R₂₁₋₂ is as defined above, (F-11)
 (c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2-
 and/or 6-, and 5- and/or 6- position with R₂₁₋₂ is as defined above, (F-12)
 30 (d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1
 or 2 R₂₁₋₂ as is defined above, (F-13)
 (e) pyrazin-2-yl optionally substituted with 1 or 2 R₂₁₋₂ as is defined
 above, (F-14)
 (f) imidazol-2-yl optionally substituted in the 1 position with C₁-C₃
 35 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2
 R₂₁₋₂ as defined above, (F-15)

- (g) 1,3,4-triazol-2-yl optionally substituted in the 1 position with C_1 - C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)
- (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1 - C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl], (F-22)
- (n) 1,2,4-triazin-3-yl optionally substituted at the 5- and/or 6- position with R_{M-2} as is defined above, (F-23)
- (2) $-(CH_2)_{2-4}$ -(1-piperazinyl) optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)
- (3) -Heteroaryl, as defined above,
- (4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is
- (a) $-O-CH_2CH_2-Y$, where Y is C_1 - C_3 alkylamino, di(C_1 - C_3)alkylamino where the alkyl groups are the same or different, C_3 - C_6 alkyleneimino, optionally substituted with 1 or 2 C_1 - C_3 alkyl,
- (b) $-NR_{21-5}CH_2CH_2-Y$, where R_{21-5} is -H or C_1 - C_3 alkyl and Y is as defined above,
- (c) $-(CH_2)_g-N(R_{21-5})$ -Heteroaryl, where g is 2, 3 or 4, and where R_{21-5} and Heteroaryl are as defined above,
- (5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1 - C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3 - C_6 heterocyclic ring and where m is as defined above,
- (6) $-(CHCH_3)_b-(CH_2)_f$ -Aryl where b is 0 and f is 1 through 4 or b is 1 and f is 0 through 3, where Aryl is as defined above,
- (7) $-(CH_2)_i$ -Heteroaryl, where i is 1 through 4 and Heteroaryl is as defined above,
- (8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-25)
- (9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by

-Heteroaryl where Heteroaryl is as defined above,

(F-26)

(B) $R_{21-\beta}$ is

(1) -H,

(2) C_1-C_3 alkyl,

5

(3) C_5-C_7 cycloalkyl,

(4) $-(CH_2)_m-NR_{21-1}$ -Heteroaryl, where m, R_{21-1} and Heteroaryl are as defined above,

(5) (1-piperazinyl)- (C_2-C_4) alkyl optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above,

(F-24)

10

(6) $-(CH_2)_m-X_4$, where m and X_4 are as defined above,

(7) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where m, R_{21-22} and R_{21-23} are as defined above,

15

(8) $-(CHCH_3)_b-(CH_2)_f-R_{21-24}$, where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1-C_3 alkoxy, $-NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1-C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4-C_7 cyclicamino ring and where b and f are as defined above,

(9) 2-pyridinylmethyl,

(C) $R_{21-\alpha}$ and $R_{21-\beta}$ are taken together with the attached nitrogen atom to form a heterocyclic ring selected from the group consisting of

20

(1) 2-(carboxy)-1-pyrrolidinyl optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt,

(F-27)

(2) 2-(carboxy)-1-piperidinyl optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt,

(F-28)

25

(3) 2-(carboxy)-1-hexamethyleneimino optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt,

(F-29)

(4) 2-(carboxy)-1-heptamethyleneimino optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt,

(F-30)

30

(5) 1-piperazinyl optionally substituted in the 4- position with $R_{21-28}CO-(CH_2)_j$ - where R_{21-28} is -Aryl, -Heteroaryl, $-NR_{21-29}$ Heteroaryl and 2-furanyl, where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 through 3, and Aryl and Heteroaryl are as defined above,

(F-31)

(6) 1-piperazinyl substituted in the 4- position with Heteroaryl- $(CH_2)_j$ -, where Heteroaryl and j are as defined above,

(F-32)

35

(7) 1-piperazinyl substituted in the 4- position with Aryl- $(CH_2)_j$ -, where Aryl and j are as defined above,

(F-33)

(8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as defined

above,

(F-34)

(9) 1-piperazinyl substituted in the 4- position with Heteroaryl-

$\text{NR}_{21-29}\text{-CO-(CH}_2\text{)}_i\text{-}$ where Heteroaryl, R_{21-29} and i are as defined above;

(F-35)

(10) 1-piperazinyl substituted in the 4- position with

5 $\text{-(CH}_2\text{)}_j\text{-C}^*=\text{C(2-pyridinyl)-N=N-C(2-pyridinyl)=C}^*\text{H}$, where $*$ and j are as defined above,

(F-36)

(11) 1-piperazinyl substituted in the 4- position with


$\text{-(CH}_2\text{)}_i\text{[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazine]}$

(F-37)

and pharmaceutically acceptable salts thereof.

10

Further disclosed is a method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing bicyclic amine of formula (III)



(III)

15

where:

20

W_2 is -O- , -S- , $\text{-NR}_{54}\text{-}$ where R_{54} is -H or $\text{C}_1\text{-C}_3$ alkyl,

n_6 is 0, 1 or 2,

R_7 is -H , $\text{C}_1\text{-C}_4$ alkyl, $\text{-CO-(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-CO-}\phi$ or -prodrug where prodrug is

$\text{PO}_2\text{-O}^-\text{ cation}^+$ where cation^+ is sodium, potassium or trialkylammonium where alkyl is $\text{C}_1\text{-C}_3$,

25

$\text{-CO-CH}_2\text{-CO-NH-CH}_2\text{-SO}_2\text{-O}^-\text{ cation}^+$ where cation^+ is as defined above,

$\text{-CO-(CH}_2\text{)}_{n_{21}}\text{-R}_{51}$ where n_{21} is 1-7 and R_{51} is -COO^- , $\text{-NR}_{51-1}\text{R}_{51-2}$ where

R_{51-1} and R_{51-2} are the same or different and are -H or $\text{C}_1\text{-C}_3$ alkyl,

$\text{-N}^+\text{R}_{51-1}\text{R}_{51-2}\text{R}_{51-3}\text{ halide}^-$ where $\text{R}_{51-1}\text{R}_{51-2}\text{R}_{51-3}$ are the same or different and are -H or $\text{C}_1\text{-C}_3$ alkyl, and where halide is -Cl or -Br ,

30

$\text{-CO-CH=CH-CO-O}^-\text{ cation}^+$ where cation^+ is as defined above,

$\text{-CO-N}^*\text{-CH=CH-N=C}^*\text{H}$ where the atoms marked with an asterisk (*) are

bonded to each other resulting in the formation of a ring,


$\text{-CO-C}^*=\text{C}[(\text{CH}_2)_{n_{22}}\text{-NH}_2]\text{-CH=CH-CH=C}^*\text{H}$ where n_{22} is 1 or 2 and where

the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a


35 ring,

$\text{-CO-C}^*=\text{CH-CH=C(-NR}_{52}\text{)-CH=C}^*\text{H}$ where R_{52} is -H or $\text{C}_1\text{-C}_3$ alkyl and

where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring,



5 C_1-C_{18} ,



above,



above,

10 $-\text{CO}-\text{NH}-\text{C}_6\text{H}_4-\text{R}_{55}$ where R_{55} is $-\text{H}$ or C_1-C_3 alkyl, $-\text{NO}_2$,

$-\text{NR}_{51-1}\text{R}_{51-2}$ where R_{51-1} and R_{51-2} are as defined above and

R_{10} is $-\text{H}$ or $-\text{CH}_3$,

R_{11} is $-\text{H}$ or $-\text{CH}_3$,

R_{12} is $-\text{H}$ or $-\text{CH}_3$,

15 (18-1) R_{16} is $\alpha\text{-R}_{16-1}:\beta\text{-R}_{16-2}$ where one of R_{16-1} and R_{16-2} is $-\text{H}$, $-\text{CH}_3$,

$-\text{CH}_2\text{CH}_3$ or $-\phi$ and the other is $-\text{X}_3-\text{NR}_{21-\alpha}\text{R}_{21-\beta}$ where X_3 is $-\text{CO}-$, $-(\text{CH}_2)_{n_{16}}-\text{CO}-$ where n_{16} is 1 or 2, $-(\text{CH}_2)_{n_3}-$ where n_3 is 1-6, or $-\text{CO}-\text{O}-(\text{CH}_2)_{n_{15}}-$ where n_{15} is 2-6, R_{25} and R_{26} are $-\text{H}:-\text{H}$;

(A) $\text{R}_{21-\alpha}$ is

20 (1) $-(\text{CH}_2)_m-\text{NR}_{21-1}\text{-Heteroaryl}$, where m is 2, 3 or 4, where R_{21-1} is $-\text{H}$ or C_1-C_3 alkyl, where Heteroaryl is:

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof

optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is

(i) $-\text{F}$,

25 (ii) $-\text{Cl}$,

(iii) $-\text{Br}$,

(iv) C_1-C_5 alkyl,

(v) $-\text{CH}_2-\text{CH}=\text{CH}_2$,

(vi) $-\text{Aryl}$, where Aryl is phenyl optionally substituted with 1

30 thru 2 $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, C_1-C_3 alkoxy, $-\text{COOH}$, $-\text{NH}_2$, C_1-C_3 alkylamino, $\text{di}(\text{C}_1-\text{C}_3)\text{alkylamino}$, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2-C_4 acylamino, $-\text{NH}-\text{CHO}$, with 1 $-\text{F}$ or $-\text{CF}_3$ or with 3,4-methylenedioxy and 3,4-ethylenedioxy;

(vii) $-\text{NR}_{21-3}\text{R}_{21-3}$ where the R_{21-3} 's are the same or different

35 and are $-\text{H}$, C_1-C_3 alkyl or $-\text{CH}_2-\text{CH}=\text{CH}_2$,

(viii) $^*\text{CH}_2-(\text{CH}_2)_q-\text{CH}_2-\text{N}^*$ where the atoms marked with an

- asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,
 (viii) $^*\text{CH}_2-(\text{CH}_2)_c-\text{G}-(\text{CH}_2)_d-\text{CH}_2-\text{N}^*$ where the atoms
 marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4),
 where G is -O-, -S-, -SO-, -SO₂- or -NR₂₁₋₄, where R₂₁₋₄ is -H, C₁-C₃ alkyl, or Aryl as
 5 defined above, where c and d are the same or different and are 0 thru 2 with the proviso that
 the total number of ring carbon atoms is 4, 5 or 6,
 (ix) 3-pyrrolin-1-yl, (F-5)
 (x) pyrrol-1-yl optionally substituted with (F-6)
 C₁-C₃ alkyl,
 10 (xi) piperidin-1-yl optionally substituted with 1 or 2 (F-7)
 C₁-C₃ alkyl, (F-8)
 (xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-9)
 (xiii) 1-hexamethyleneimino containing a 3- or 4- double bond
 or 3- and 5- double bonds, (F-10)
 15 (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by
 two C₁-C₃ alkyl being the same or different, (F-10)
 (xv) -OH,
 (xvi) C₁-C₃ alkoxy,
 (xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where
 20 R₂₁₋₇ is -H or C₁-C₃ alkyl and e is 0 thru 3,
 (xviii) pyridin-2-, 3- or 4-yl,
 (xix) -CF₃,
 (xx) -CCl₃,
 (xxi) -SCH₃,
 25 (b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the
 4- and/or 6- position with R₂₁₋₂ is as defined above, (F-11)
 (c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2-
 and/or 6-, and 5- and/or 6- position with R₂₁₋₂ is as defined
 above, (F-12)
 30 (d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1
 or 2 R₂₁₋₂ as is defined above, (F-13)
 (e) pyrazin-2-yl optionally substituted with 1 or 2 R₂₁₋₂ as is defined
 above, (F-14)
 (f) imidazol-2-yl optionally substituted in the 1 position with C₁-C₃
 35 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2
 R₂₁₋₂ as defined above, (F-15)

- (g) 1,3,4-triazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)
- (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl], (F-22)
- (n) 1,2,4-triazin-3-yl optionally substituted at the 5- and/or 6- position with R_{21-2} as is defined above, (F-23)
- (2) $-(CH_2)_{2-4}$ -(1-piperazinyl) optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)
- (3) -Heteroaryl, as defined above,
- (4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is
- (a) $-O-CH_2CH_2-Y$, where Y is C_1-C_3 alkylamino, di(C_1-C_3)alkylamino where the alkyl groups are the same or different, C_3-C_6 alkyleneimino, optionally substituted with 1 or 2 C_1-C_3 alkyl,
- (b) $-NR_{21-20}CH_2CH_2-Y$, where R_{21-20} is -H or C_1-C_3 alkyl and Y is as defined above,
- (c) $-(CH_2)_g-N(R_{21-20})$ -Heteroaryl, where g is 2, 3 or 4, and where R_{21-20} and Heteroaryl are as defined above,
- (5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1-C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3-C_6 heterocyclic ring and where m is as defined above,
- (6) $-(CHCH_3)_b-(CH_2)_f$ -Aryl where b is 0 and f is 1 thru 4 or b is 1 and f is 0 thru 3, where Aryl is as defined above,
- (7) $-(CH_2)_i$ -Heteroaryl, where i is 1 thru 4 and Heteroaryl is as defined above,
- (8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-25)
- (9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by

-Heteroaryl where Heteroaryl is as defined above, and

(F-26)

(B) $R_{21-\beta}$ is

(1) -H.

(2) C_1-C_3 alkyl,

(3) C_5-C_7 cycloalkyl,

(4) $-(CH_2)_m-NR_{21-1}$ -Heteroaryl, where m, R_{21-1} and Heteroaryl are as defined

above,

(5) (1-piperazinyl)-(C_2-C_4)alkyl optionally substituted in the 4- position with -

Aryl or -Heteroaryl as defined above,

(F-24)

(6) $-(CH_2)_m-X_4$, where m and X_4 are as defined above,

(7) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where m, R_{21-22} and R_{21-23} are as defined

above,

(8) $-(CHCH_3)_b-(CH_2)_fR_{21-24}$, where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1-C_3 alkoxy, $-NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1-C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4-C_7 cyclicamino ring and where b and f are as defined above,

(9) 2-pyridinylmethyl,

(10) 2-phenylethyl,

(C) $R_{21-\alpha}$ and $R_{21-\beta}$ are taken together with the attached nitrogen atom to form a

heterocyclic ring selected from the group consisting of

(1) 2-(carboxy)-1-pyrrolidinyl optionally as the C_1-C_3 alkyl ester or as a

pharmaceutically acceptable salt,

(F-27)

(2) 2-(carboxy)-1-piperidinyl optionally as the C_1-C_3 alkyl ester or as a

pharmaceutically acceptable salt,

(F-28)

(3) 2-(carboxy)-1-hexamethyleneimino optionally as the C_1-C_3 alkyl ester or as

a pharmaceutically acceptable salt,

(F-29)

(4) 2-(carboxy)-1-heptamethyleneimino optionally as the C_1-C_3 alkyl ester or as

a pharmaceutically acceptable salt,

(F-30)

(5) 1-piperazinyl optionally substituted in the 4- position with $R_{21-28}CO-$

$(CH_2)_j-$ where R_{21-28} is -Aryl, -Heteroaryl, $-NR_{21-29}$ -Heteroaryl and 2-furanyl, where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 thru 3, and Aryl and Heteroaryl are as defined above, (F-31)

(6) 1-piperazinyl substituted in the 4-position with Heteroaryl $(CH_2)_j-$ where

Heteroaryl and j are as defined above

(F-32)

(7) 1-piperazinyl substituted in the 4-position with Aryl- $(CH_2)_j-$, where Aryl

and j are as defined above,

(F-33)

(8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as defined

above,

(F-34)

(9) 1-piperazinyl substituted in the 4- position with Heteroaryl-NR₂₁₋₂₉-CO-(CH₂)_i, where Heteroaryl, R₂₁₋₂₉ and i are as defined above;

(F-35)

(10) 1-piperazinyl substituted in the 4- position with

5 -(CH₂)_j-C^{*}=C(2-pyridinyl)-N=N-C(2-pyridinyl)=C^{*}H, where * and j are as defined above,

(F-36)

(11) 1-piperazinyl substituted in the 4- position with -(CH₂)_i-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazine] where i is as defined

above,

(F-37)


10 (12) 1-piperazinyl substituted in the 4- position with C₁-C₃ alkyl optionally substituted with 1 or 2 Aryl;

(18-2) n₆ is 0, R₁₆ is R₁₆₋₃:R₁₆₋₄ and R₂₅ is R₂₅₋₃:R₂₅₋₄ where one of R₁₆₋₃ and R₁₆₋₄ is taken together with one of R₂₅₋₃ and R₂₅₋₄ to form a second bond between the carbon atoms to which R₁₆ and R₂₅ are attached and the other of R₁₆₋₃ and R₁₆₋₄ is -X₃-NR_{21-α}R_{21-β}

15 where X₃, R_{21-α} and R_{21-β} are as defined above and the other of R₂₅₋₃ and R₂₅₋₄ is -H,

(18-3) n₆ is 1, R₂₅ is R₂₅₋₅ and R₂₅₋₆ and R₂₆ is R₂₆₋₅ and R₂₆₋₆ where one of R₂₅₋₅ and R₂₅₋₆ and one of R₂₆₋₅ and R₂₆₋₆ are taken together to form a second bond between the carbon atoms to which R₂₅ and R₂₆ are attached and the other of R₂₅₋₅ and R₂₅₋₆ and R₂₆₋₅ and R₂₆₋₆ are -H, and pharmaceutically acceptable salts thereof.

20 Additionally disclosed is a method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing bicyclic ether of formula (IV)



(IV)

25

where R₁ is -H or -OR₁₋₁ where R₁₋₁ is C₁-C₃ alkyl and where R₂ is -H or -OR₂₋₁ where R₂₋₁ is C₁-C₃ alkyl with the proviso that R₂ is not -H only when R₁ is -H,

30 R₃ is α-R₃₋₁:β-R₃₋₂ where R₃₋₁ and R₃₋₂ are the same or different and are -H or -CH₃ with the proviso that R₃₋₂ is not -CH₃ unless R₃₋₁ is -CH₃,

n is 1, 2 or 3,

R₄ is R₄₋₁:R₄₋₂ where R₄₋₁ is -H, -CH₃, -CH₂CH₃, 4-fluorophenyl, 4-chlorophenyl, R₄₋₂ is -(CH₂)_n-R₄₋₃ where n is 1, 2 or 3 and where R₄₋₃ is

35

-Cl,

1-piperazinyl optionally substituted in the 4-position with a member selected

from the group consisting of

- ϕ optionally substituted with 1 -CF₃, -Cl, -F, -CH₃, -CH₂CH₃,

2-pyridinyl optionally substituted in the 6-position with -NR₄₋₄R₄₋₅

where R₄₋₄ and R₄₋₅ are the same or different and are -H, C₁-C₃ alkyl and where R₄₋₄ and R₄₋₅

5 are taken together with the attached nitrogen atom to form a ring selected from the group consisting of 1-pyrrolidinyl, 1-piperidinyl, 1-morpholinyl,

4-pyrimidinyl optionally substituted in the 2 and/or 6-position with -

NR₄₋₄R₄₋₅ where R₄₋₄ and R₄₋₅ are as defined above,

piperid-3-en-1-yl optionally substituted in the 4-position with a

10 member selected from the group consisting of

- ϕ optionally substituted with 1 -CF₃, -Cl, -F, -CH₃, -CH₂CH₃,

2-pyridinyl optionally substituted in the 6-position with -NR₄₋₄R₄₋₅

where R₄₋₄ and R₄₋₅ are the same or different and are -H, C₁-C₃alkyl and where R₄₋₄ and R₄₋₅

are taken together with the attached nitrogen atom to form a ring selected from the group

15 consisting of 1-pyrrolidinyl, 1-piperidinyl, 1-morpholinyl,


4-pyrimidinyl optionally substituted in the 2 and/or 6-position with -

NR₄₋₄R₄₋₅ where R₄₋₄ and R₄₋₅ are as defined above, and pharmaceutically acceptable salts thereof.

Further disclosed is a method of treating resistance to cancer chemotherapeutic agents

20 in human cancer patients which comprises administering to that human an effective amount of a sensitizing tricyclic compound of formula (V)

25



(V)

30 where:

n₁ is 1 thru 3,

R₁ is α -R₁₋₁: β -R₁₋₂ where R₁₋₁ and R₁₋₂ are the same or different and are -H,

C₁-C₃ alkyl,


R₂ is C₁-C₃ alkyl,

35 R₃ is - ϕ optionally substituted with 1 thru 3 -F, -Cl, C₁-C₃ alkyl and pharmaceutically acceptable salts thereof.

Also disclosed are indoles of formula (VI)

5

10



and a method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing indole of

15 formula (VI) where:

R_1 is $-C=N$ or $-CO-NH_2$;

n_1 is 1 thru 5;

R_{21} is

(1) $R_{21-28}-CO-(CH_2)_j-$ where R_{21-28} is -Aryl, $-NR_{21-29}$ Aryl and 2-furanyl,
 20 where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 thru 3 and Aryl is phenyl optionally substituted with 1 or 2 -Cl, -Br, C_1-C_3 alkoxy, $-COOH$, $-NH_2$, C_1-C_3 alkylamino, di(C_1-C_3)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2-C_4 acylamino and $-NH-CHO$ or with 1 -F or $-CF_3$;

(F-31)

25 (2) Heteroaryl- $(CH_2)_j-$, where Heteroaryl is

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is

(i) -F,

(ii) -Cl,

30

(iii) -Br,

(iv) C_1-C_5 alkyl,

(v) $-CH_2-CH=CH_2$,

(vi) -Aryl, where Aryl is as defined above,

(vii) $-NR_{21-3}R_{21-3}$ where the R_{21-3} 's are the same or different

35 and are -H, C_1-C_3 alkyl or $-CH_2-CH=CH_2$,

(viii) $^*CH_2-(CH_2)_q-CH_2-N^*$ - where the atoms marked with an

- asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,
 (viii) $^*\text{CH}_2\text{-CH}_2\text{-(CH}_2\text{)}_c\text{-G-(CH}_2\text{)}_d\text{-CH}_2\text{-CH}_2\text{-N}^*$ where the
 atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring
 (F-4), where G is -O-, -S-, -SO-, -SO₂- or -NHR₂₁₋₄, where R₂₁₋₄ is -H, C₁-C₃ alkyl, or Aryl
 5 as defined above, where c and d are the same or different and are 0 thru 2 with the proviso that
 the total number of ring carbon atoms is 4, 5 or 6,
 (ix) 3-pyrrolin-1-yl, (F-5)
 (x) pyrrol-1-yl optionally substituted with C₁-C₃ alkyl, (F-6)
 (xi) piperidin-1-yl optionally substituted with 1 or 2 C₁-C₃
 10 alkyl, (F-7)
 (xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)
 (xiii) 1-hexamethyleneimino containing a 3- or 4- double bond
 or 3- and 5- double bonds, (F-9)
 (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by
 15 two C₁-C₃ alkyl being the same or different, (F-10)
 (xv) -OH,
 (xvi) C₁-C₃ alkoxy,
 (xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where R₂₁₋₇
 is -H or C₁-C₃ alkyl and e is 0 thru 3,
 20 (xviii) pyridin-2-, 3- or 4-yl,
 (b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the
 4- and/or 6- position with R₂₁₋₂ is as defined above, (F-11)
 (c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2-
 and/or 6- position with R₂₁₋₂ is as defined above, (F-12)
 25 (d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1
 or 2 R₂₁₋₂ as is defined above, (F-13)
 (e) pyrazin-2-yl optionally substituted with 1 or 2 R₂₁₋₂ as is defined
 above, (F-14)
 (f) imidazol-2-yl optionally substituted in the 1 position with C₁-C₃
 30 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2
 R₂₁₋₂ as defined above, (F-15)
 (g) 1,2,4-triazol-3-yl optionally substituted in the 1 position with C₁-C₃
 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R₂₁₋₂ as
 defined above, (F-16)
 35 (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C₁-
 C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or

- 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- 5 (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]-ethyl]piperazinyl, (F-22)
- (n) 1,2,4-triazol-3-yl optionally substituted at the 5- and/or 6- position with R_{21-2} as is defined above, (F-23)
- 10 and where j is as defined above, (F-32)
- (3) Aryl-(CH₂)_j-, where Aryl and j are as defined above, (F-33)
- and pharmaceutically acceptable salts thereof.

DETAILED DESCRIPTION OF THE INVENTION

The steroidal amines (I) are known, see International Publication No. WO87/01706,
15 published March 26, 1987 based on International Patent Application No. PCT/US86/01797.

The alkyl amines (II) and the bicyclic amines (III) are known, see International Publication No. WO88/08424, published November 3, 1988 based on International Patent Application No. PCT/US88/01212.

The bicyclic ethers (IV) are known, see for example, US Patents 4,206,123, 4,577,021,
20 4,711,960 and 4,487,774.

The tricyclic amines (V) are known see US Patent 4,487,774.

The indole amines (VI) are prepared by the process set forth in EXAMPLES 5-8.

With either the steroidal amines (I), alkyl amines (II) or bicyclic amines (III), it is preferred that the amine portion be cyclized, that is $R_{21-\alpha}$ and $R_{21-\beta}$ be taken together with the
25 attached nitrogen atom to form a heterocyclic ring which is group (C). It is further preferred that the heterocyclic ring be piperazinyl substituted with with either Aryl or Heteroaryl, preferably with Heteroaryl. It is preferred that the Heteroaryl substituent itself be substituted, more preferably be di substituted. It is preferred that the substituents on the Heteroaryl group themselves be cyclized such as pyrrolidinyl.

30 It is preferred that the steroidal amine (I) be
17 α -hydroxy-21-[4-(2-pyridinyl)-1-piperazinyl]-pregna-4,9(11)-diene-3,20-dione,
21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-
triene-3,20-dione,
21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]pregna-1,4,9(11)-triene-3,20-
35 dione,
21-[4-3,6-bis(diethylamino)-2-pyridinyl]-1-piperazinyl-16 α -methylpregna-1,4,9(11)-

triene-3,20-dione,

21-[4-(4,6-di-1-pyrrolidinyl-1,3,5-triazin-2-yl)-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

21-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

21-[4-(4,6-di-1-pyrrolidinyl-2-pyrimidinyl)-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione. It is more preferred that the steroidal amine (I) be

17 α -hydroxy-21-[4-(2-pyridinyl)-1-piperazinyl]-pregna-4,9(11)-diene-3,20-dione,

21-[4-2,6-di-1-pyrrolidinyl-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]pregna-1,4,9(11)-triene-3,20-dione,

21-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione. It is even more preferred that the steroidal amine be 21-[4-2,6-di-1-pyrrolidinyl]-4-

pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione.

It is preferred that the alkyl amines (II) be

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinehexanol,

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazineoctanol,

4-[[6-[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]hexyl]oxy]phenol,

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)- α -phenyl-1-piperazinebutanol,

4-[3-[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]propyl]-2,5-dimethylphenol,

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazineheptanoic acid methyl ester. It is more preferred that the alkyl amines (II) be

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazineoctanol,

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)- α -phenyl-1-piperazinebutanol.

It is preferred that the bicyclic amines (III) be

2-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]methyl]-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol,

2-[[4-3-(ethylamino)-2-pyridinyl]-1-piperazinyl]methyl]-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol,

3-[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]-6-(acetyloxy)-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-2-carboxylic acid propyl ester,

1-[3-(ethylamino)-2-pyridinyl]-4-[(5-methoxy-4,6,7-trimethyl-1H-indol-2-yl)carbonyl]-piperazine. It is more preferred that the bicyclic amine (III) be

2-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]methyl]-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol,

2-[[4-3-(ethylamino)-2-pyridinyl]-1-piperazinyl)methyl]-3,4-dihydro-2,5,7,8-tetra-
methyl-2H-1-benzopyran-6-ol.

It is preferred that the bicyclic ethers (IV) be

- 1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-[3-(trifluoromethyl)-
5 phenyl]piperazine,
1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)-
piperazine,
1-[(3,4-dihydro-8-methoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)piperazine,
4-(4-chlorophenyl)-1-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-
10 1,2,3,6-tetrahydropiperidine,
1-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)-
piperazine,
1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-
yl)ethyl]piperazine,
15 1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(4-
fluorophenyl)piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-[3-
(trifluoromethyl)phenyl]piperazine,
1-(4-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-
20 yl)ethyl]piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-
methylphenyl)piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,6-
tetrahydro-4-phenylpyridine,
25 2-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,5-
tetrahydro-6,7-dimethoxyisoquinoline,
1-[2-(3,4-dihydro-5,6-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-
phenylpiperazine,
1-(4-fluorophenyl)-4-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-
30 benzopyran-1-yl]-propyl]piperazine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-(2-
methylphenyl)piperazine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-
phenylpiperazine,
35 2H-Benzimidazol-2-one, 1-[1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-
benzopyran-1-yl]-propyl]-4-piperidinyl]-1,3-dihydropiperazine,

- 1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-(2-pyridinyl)piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,6-tetrahydro-4-phenylpyridine,
5 1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,
1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]piperazine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-
10 phenylpiperazine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-(2-methylphenyl)piperazine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-(4-fluorophenyl)piperazine,
15 1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl]propyl]-4-phenylpiperazine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4-dimethyl-1H-2-benzopyran-1-yl)propyl]-1,2,3,6-
20 tetrahydro-4-phenylpyridine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-4-phenylpiperazine,
1-(3-chloropropyl)-1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran,
25 1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4-dimethyl-1H-2-benzopyran-1-yl)propyl]-4-(2-methylphenyl)piperazine,
1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-1-methyl-1H-2-benzopyran-1-
30 yl)ethyl]piperazine,
1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]piperazine,
1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]piperazine,
35 6-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-N,N,N',N'-tetraethyl-2,4-pyrimidinediimine,

4-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-2,6-di-1-pyrrolidinylpyrimidine. It is more preferred that the bicyclic ethers (IV) be

1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)-piperazine,

5 1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,

1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,

1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,

1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]piperazine,

4-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-2,6-di-1-pyrrolidinylpyrimidine. It is most preferred that the bicyclic ethers (IV) be 1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine.

It is preferred that the tricyclic amines (V) be

1-(4-fluorophenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine,

1-(2-methylphenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine,

1-(2-chlorophenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine. It is more preferred that the tricyclic amines (V) be 1-(2-chlorophenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine.

25 It is preferred that the indole amines (VI) be

4-[3-4-diphenylmethyl]-1-piperazinyl]propoxy]indole-2-carboxamide,

4-[3-[4-[2,4-dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indol-2-carboxamide,

2-cyano-4-[3-[4-(diphenylmethyl)-1-piperazinyl]propoxy]indole,

2-cyano-4-[3-[4-(2,4-dipyrrolidino-6-pyrimidinyl)-1-piperazinyl]propoxy]indole. It is more preferred that the indole amines (VI) be 4-[3-4-diphenylmethyl]-1-piperazinyl]propoxy]indole-2-carboxamide.

Other compounds which are useful in the present invention are

11-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]acetyl]oxy]-16 α -methylpregn-4-en-3,20-dione,

35 4-[5-(benzoyloxy)-2,6-di-1-pyrrolidinyl-4-pyrimidinyl]1-piperazineheptanoic acid methyl ester,

21-[4-(2,6-di-1-pyrrolidinyl-5-(4-chlorobenzoyloxy))-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

3-[2-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]ethyl]octahydro-7-[2-(5-hydroxy-2-methylphenyl)ethyl]-3 α -methyl-5H-inden-5-one.

5 It is preferred that the sensitizing compound be selected from the group consisting of 1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)-piperazine,

1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,

10 1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,

1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,

15 1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]piperazine,

11-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]acetyl]oxy]-16 α -methylpregn-4-en-3,20-dione,

21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

20 4-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-2,6-di-1-pyrrolidinylpyrimidine,

21-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)- α -phenyl-1-piperazinebutanol,

25 4-[3-4-diphenylmethyl]-1-piperazinyl]propoxy]indole-2-carboxamide,

4-[5-(benzoyloxy)-2,6-di-1-pyrrolidinyl-4-pyrimidinyl]1-piperazineheptanoic acid methyl

ester.

The sensitizing compounds of this invention are amines and as such form acid addition salts when reacted with acids of sufficient strength. Pharmaceutically acceptable salts include salts of both inorganic and organic acids. The pharmaceutically acceptable salts are preferred over the corresponding free amines since they produce compounds which are more water soluble and more crystalline. The preferred pharmaceutically acceptable salts include salts of the following acids methanesulfonic, hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, maleic, CH₃-(CH₂)_n-COOH where n is 0 thru 4, HOOC-(CH₂)_n-COOH where n is as defined above.

35 Further, the sensitizing compounds and salts thereof form solvates and hydrates thereof,

as is known in the art, which are equivalent to the nonsolvated or nonhydrated forms.

The term "treating" as used in this invention means both (1) preventing resistance to the chemotherapeutic agents and (2) overcoming resistance to the chemotherapeutic agents which already exists.

- 5 Further, "preventing" means to prevent all-together or to slow the pace at which resistance develops.

The method of the present invention is useful in sensitizing desensitized cancer cells of the following types of cancer: ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla
10 of Vater, multiple myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma. It is preferred that the cancer cells be breast cancer, multiple myeloma, ovarian or lung.

- It is realized that the desensitized cancer cells may be desensitized to more than one chemotherapeutic agent. If so, the method of the present invention will sensitize the
15 desensitized cancer cells to most of the chemotherapeutic agents to which they are desensitized.

The chemotherapeutic agents to which the cancer cells become desensitized are selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

- 20 [S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,

(S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[[[(phenylamino)carbonyl]-opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-benzofurancarboxamide,

- 25 (7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide. It is preferred that the chemotherapeutic agent be selected from the group consisting of doxorubicin or vincristine.

- 30 It is realized that new chemotherapeutic agents against cancer will be developed after this invention. The new chemotherapeutic agents to which resistance develops and which can be treated by the method of this invention are equivalent to those set forth in this invention.

- The sensitizing compounds of the present invention are used as discussed below. Patients with cancer which once responded to a chemotherapeutic agent(s) and which now
35 responds to the chemotherapeutic agent in a much poorer way, are pre-treated with the sensitizing compound for about 12-36 hours, preferably about 24 hrs, before treatment with

the desired chemotherapeutic agent(s) is resumed. Once treatment is reinstated with the chemotherapeutic agent, the sensitizing compound and chemotherapeutic agent(s) is administered concurrently. Alternatively, in individuals who have cancer and who have not been previously treated with chemotherapeutic agents, the sensitizing compounds of this invention are given when the chemotherapeutic agents are initially given to either totally prevent resistance from developing or slow the rate at which the resistance develops.

An effective amount of the sensitizing compound is from about 1 mg/kg/day to about 500 mg/kg/day, preferably from about 1 mg/kg/day to about 100 mg/kg/day, more preferably from about 5 mg/kg/day to about 75 mg/kg/day.

The sensitizing compound is administered IV, orally and IP. When administered IV it is given continuously. When it is administered orally it is given 3 or 4 times daily in divided doses.

The various sensitizing compounds of this invention need not be used separately. They can be used in combination with each other or with other known sensitizing compounds.

The exact dosage and frequency of administration depends on the particular sensitizing compound(s) used, the particular condition being treated, the severity of the condition being treated, the age, weight, general physical condition of the particular patient, other medication the individual may be taking as is well known to those skilled in the art and can be more accurately determined by measuring the blood level or concentration of the sensitizing compound(s) in the patient's blood and/or the patient's response to the particular condition being treated.

Whether or not a compound is a good sensitizing compound can readily be determined by known means, see *Cancer Letters* 50, 45-51 (1990).

DEFINITIONS AND CONVENTIONS

The definitions and explanations below are for the terms as used throughout this entire document including both the specification and the claims.

I. CONVENTIONS FOR FORMULAS AND DEFINITIONS OF VARIABLES

The chemical formulas representing various compounds or molecular fragments in the specification and claims may contain variable substituents in addition to expressly defined structural features. These variable substituents are identified by a letter or a letter followed by a numerical subscript, for example, "Z_i" or "R_i" where "i" is an integer. These variable substituents are either monovalent or bivalent, that is, they represent a group attached to the formula by one or two chemical bonds. For example, a group Z₁ would represent a bivalent variable if attached to the formula CH₃-C(=Z₁)H. Groups R_i and R_j would represent monovalent variable substituents if attached to the formula CH₃-CH₂-C(R_i)(R_j)-H. When chemical formulas are drawn in a linear fashion, such as those above, variable substituents

contained in parentheses are bonded to the atom immediately to the left of the variable substituent enclosed in parenthesis. When two or more consecutive variable substituents are enclosed in parentheses, each of the consecutive variable substituents is bonded to the immediately preceding atom to the left which is not enclosed in parentheses. Thus, in the

5 formula above, both R_i and R_j are bonded to the preceding carbon atom. Also, for any molecule with an established system of carbon atom numbering, such as steroids, these carbon atoms are designated as C_i , where "i" is the integer corresponding to the carbon atom number. For example, C_6 represents the 6 position or carbon atom number in the steroid nucleus as traditionally designated by those skilled in the art of steroid chemistry. Likewise the term " R_6 "

10 represents a variable substituent (either monovalent or bivalent) at the C_6 position.

Chemical formulas or portions thereof drawn in a linear fashion represent atoms in a linear chain. The symbol "-" in general represents a bond between two atoms in the chain. Thus $CH_3-O-CH_2-CH(R_i)-CH_3$ represents a 2-substituted-1-methoxypropane compound. In a similar fashion, the symbol "=" represents a double bond, e.g., $CH_2=C(R_i)-O-CH_3$, and the

15 symbol "≡" represents a triple bond, e.g., $HC\equiv C-CH(R_i)-CH_2-CH_3$. Carbonyl groups are represented in either one of two ways: -CO- or -C(=O)-, with the former being preferred for simplicity.

Chemical formulas of cyclic (ring) compounds or molecular fragments can be represented in a linear fashion. Thus, the compound 4-chloro-2-methylpyridine can be

20 represented in linear fashion by $N^*=C(CH_3)-CH=CCl-CH=C^*H$ with the convention that the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring. Likewise, the cyclic molecular fragment, 4-(ethyl)-1-piperazinyll can be represented by $-N^*-(CH_2)_2-N(C_2H_5)-CH_2-C^*H_2$.

A rigid cyclic (ring) structure for any compounds herein defines an orientation with

25 respect to the plane of the ring for substituents attached to each carbon atom of the rigid cyclic compound. For saturated compounds which have two substituents attached to a carbon atom which is part of a cyclic system, $-C(X_1)(X_2)-$ the two substituents may be in either an axial or equatorial position relative to the ring and may change between axial/equatorial. However, the position of the two substituents relative to the ring and each other remains fixed. While either

30 substituent at times may lie in the plane of the ring (equatorial) rather than above or below the plane (axial), one substituent is always above the other. In chemical structural formulas depicting such compounds, a substituent (X_1) which is "below" another substituent (X_2) will be identified as being in the alpha (α) configuration and is identified by a broken, dashed or dotted line attachment to the carbon atom, i.e., by the symbol "- - -" or "...". The corresponding

35 substituent attached "above" (X_2) the other (X_1) is identified as being in the beta (β) configuration and is indicated by an unbroken line attachment to the carbon atom.

When a variable substituent is bivalent, the valences may be taken together or separately or both in the definition of the variable. For example, a variable R_i attached to a carbon atom as $-C(=R_i)-$ might be bivalent and be defined as oxo or keto (thus forming a carbonyl group $(-CO-)$ or as two separately attached monovalent variable substituents $\alpha-R_{i-j}$ and $\beta-R_{i-k}$. When a bivalent variable, R_i , is defined to consist of two monovalent variable substituents, the convention used to define the bivalent variable is of the form " $\alpha-R_{i-j}:\beta-R_{i-k}$ " or some variant thereof. In such a case both $\alpha-R_{i-j}$ and $\beta-R_{i-k}$ are attached to the carbon atom to give $-C(\alpha-R_{i-j})(\beta-R_{i-k})-$. For example, when the bivalent variable R_6 , $-C(=R_6)-$ is defined to consist of two monovalent variable substituents, the two monovalent variable substituents are $\alpha-R_{6-1}:\beta-R_{6-2}$, $\alpha-R_{6-9}:\beta-R_{6-10}$, etc, giving $-C(\alpha-R_{6-1})(\beta-R_{6-2})-$, $-C(\alpha-R_{6-9})(\beta-R_{6-10})-$, etc. Likewise, for the bivalent variable R_{11} , $-C(=R_{11})-$, two monovalent variable substituents are $\alpha-R_{11-1}:\beta-R_{11-2}$. For a ring substituent for which separate α and β orientations do not exist (e.g. due to the presence of a carbon carbon double bond in the ring), and for a substituent bonded to a carbon atom which is not part of a ring the above convention is still used, but the α and β designations are omitted.

Just as a bivalent variable may be defined as two separate monovalent variable substituents, two separate monovalent variable substituents may be defined to be taken together to form a bivalent variable. For example, in the formula $-C_1(R_i)H-C_2(R_j)H-$ (C_1 and C_2 define arbitrarily a first and second carbon atom, respectively) R_i and R_j may be defined to be taken together to form (1) a second bond between C_1 and C_2 or (2) a bivalent group such as oxa $(-O-)$ and the formula thereby describes an epoxide. When R_i and R_j are taken together to form a more complex entity, such as the group $-X-Y-$, then the orientation of the entity is such that C_1 in the above formula is bonded to X and C_2 is bonded to Y . Thus, by convention the designation "... R_i and R_j are taken together to form $-CH_2-CH_2-O-CO-$..." means a lactone in which the carbonyl is bonded to C_2 . However, when designated "... R_j and R_i are taken together to form $-CO-O-CH_2-CH_2-$ the convention means a lactone in which the carbonyl is bonded to C_1 .

The carbon atom content of variable substituents is indicated in one of two ways. The first method uses a prefix to the entire name of the variable such as " C_1-C_4 ", where both "1" and "4" are integers representing the minimum and maximum number of carbon atoms in the variable. The prefix is separated from the variable by a space. For example, " C_1-C_4 alkyl" represents alkyl of 1 through 4 carbon atoms, (including isomeric forms thereof unless an express indication to the contrary is given). Whenever this single prefix is given, the prefix indicates the entire carbon atom content of the variable being defined. Thus C_2-C_4 alkoxy carbonyl describes a group $CH_3-(CH_2)_n-O-CO-$ where n is zero, one or two. By the second method the carbon atom content of only each portion of the definition is indicated separately by enclosing the " C_i-C_j " designation in parentheses and placing it immediately (no

intervening space) before the portion of the definition being defined. By this optional convention (C₁-C₃)alkoxycarbonyl has the same meaning as C₂-C₄ alkoxycarbonyl because the "C₁-C₃" refers only to the carbon atom content of the alkoxy group. Similarly while both C₂-C₆ alkoxyalkyl and (C₁-C₃)alkoxy(C₁-C₃)alkyl define alkoxyalkyl groups containing from 2 to 6 carbon atoms, the two definitions differ since the former definition allows either the alkoxy or alkyl portion alone to contain 4 or 5 carbon atoms while the latter definition limits either of these groups to 3 carbon atoms.

When the claims contain a fairly complex (cyclic) substituent, at the end of the phrase naming/designating that particular substituent will be a notation in (parentheses) which will correspond to the same name/designation in one of the CHARTS which will also set forth the chemical structural formula of that particular substituent.

II. DEFINITIONS

Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance and bioavailability.

Treating means both (1) preventing resistance to the chemotherapeutic agents and (2) overcoming resistance to the chemotherapeutic agents which already exists.

Preventing means to prevent all-together or to slow the pace at which resistance develops.

Saline refers to a saturated aqueous sodium chloride solution.

TLC refers to thin-layer chromatography.

Ether refers to diethyl ether.

EXAMPLES

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, practice the present invention to its fullest extent. The following detailed examples describe how to perform the various processes of the invention and are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures.

EXAMPLE 1 Non-Hodgkin's Lymphoma Treated With Doxorubin, Vincristine, Cyclophosphamide and Dexamethasone

A 70 kg male patient with non-Hodgkin's Lymphoma was treated with doxorubin, vincristine, cyclophosphamide and dexamethasone and after a period of time the lymphoma did not respond as well as it did previously to the treatment. Therefore, to sensitize the lymphoma to further treatment with with these agents the patient is treated as follows:

- 21-[4-2,6-di-1-pyrrolidinyl-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione [IV, 0.01 mg/kg/hr to 5.0 mg/kg/hr for days 1-5],
 cyclophosphamide [IV, 600 mg/m² on day 2],
 vincristine [IV, 24 hr infusion, 0.4 mg/day, days 2-5],
 5 doxorubicin [IV, 24 hr infusion, 10 mg/m²/day, days 2-5],
 dexamethasone [orally, 40 mg/day, days 2-5].

The treatment is repeated every 3 to 4 weeks in the absence of severe systemic toxicities, see J. Clin Oncol. 9, 17 (1991).

EXAMPLE 2 Pancreatic Carcinoma Treated With Adriamycin

- 10 A 55 kg female patient with pancreatic carcinoma was treated with adriamycin and after a period of time the carcinoma did not respond as well as it did previously to the treatment. Therefore, to sensitize the carcinoma to treatment with adriamycin the patient is treated as follows:

- 4-[3-4-diphenylmethyl)-1-piperazinyl]propoxy]indole-2-carboxamide, [IV, 0.01
 15 mg/kg/hr to 5.0 mg/kg/hr for days 1-5],
 adriamycin [IV, 50 mg/m²/day for days 2-5].

The course of treatment is repeated every 3 weeks with adjustment of adriamycin dose if toxicities in blood counts or severe stomatitis are noted as is known to those skilled in the art; see Am. J. Clin. Oncol., 9, 355 (1986).

20 EXAMPLE 3 Breast Cancer Treated With Adriamycin

- A 62 kg female patient with breast cancer was treated with adriamycin and after a period of time the cancer did not respond as well as it did previously to the treatment. Therefore, to sensitize the breast cancer to treatment with adriamycin the patient is treated as follows:

- 25 11-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]acetyl]oxy]-16 α -methylpregn-4-en-3,20-dione, [IV, 50 mg/kg/day, over 15-20 min four times a day on days 1-6],
 adriamycin [IV infusion at 60 mg/day on days 2-6]

The course of treatment is repeated every 3 to 4 weeks in the absence of severe systemic toxicity; see J. Clin. Oncol., 6, 880 (1988).

30 EXAMPLE 4 Multiple Myeloma Treated with Vincristine, Doxorubicin, Dexamethasone, and Cyclophosphamide

- A 75 kg male with diagnosed multiple myeloma (who has never been treated for cancer previously) is treated as follows:

- 1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-
 35 methyphenyl)piperazine, [IV, 0.01 mg/kg/hr to 5.0 mg/kg/hr for days 1-5],
 cyclophosphamide [IV, 600 mg/m² on day 2],

vincristine [IV, 24 hr infusion, 0.4 mg/day, days 2-5],
doxorubicin [IV, 24 hr infusion, 10 mg/m²/day, days 2-5],
dexamethasone [orally, 40 mg/day, days 2-5].

The treatment is repeated every 3 to 4 weeks in the absence of severe systemic toxicities, see J.

5 Clin Oncol. 9, 17 (1991).

EXAMPLE 5 4-[3-[4-(Diphenylmethyl)-1-piperazinyl] propoxy]indole-2-carboxamide
(VI)

4-(Benzyloxy)indole-2-carboxylic acid [mp 239-241° d., lit. mp 241-242° d.] (15.0 g)
in 250 ml of methylene chloride and 3 ml of DMF is treated dropwise, under nitrogen, with
10 4.2 ml (6.85 g) of thionyl chloride in 65 ml of methylene chloride over 10 min. The mixture
is stirred at 20-25° for 2.5 hr, 250 ml of ether is added, a -20° bath was applied, and ammonia
(gas) is bubbled through the mixture for 30 min. The mixture is stirred as the bath temperature
was allowed to rise from -20° to 10° over 1 hr. After another hour at 20-25° the mixture is
bubbled with a stream of nitrogen for 30 min then concentrated under reduced pressure. The
15 residue is partitioned between 150 ml of water and 600 ml of methylene chloride (a solid
formed at the interface). The aqueous layer (and solid) is extracted with additional methylene
chloride (2 x 600 ml); the solid persisted in the aqueous layer. This is extracted with ethyl
acetate (2 x 400 ml). The pooled methylene chloride extract is washed with saline, dried over
magnesium sulfate and concentrated. The pooled ethyl acetate extract was dried over
20 magnesium sulfate and concentrated. After crystallization from ethanol/water and recrystalliz-
ation from ethanol an analytical sample had mp 188-9°. The proposed structure is supported
by NMR, IR and mass spectra as 4-(benzyloxy)indole-2-carboxamide.

4-(benzyloxy)indole-2-carboxamide (4.98 g) in 300 ml of methanol is stirred under 1
atm of hydrogen with 0.75 g of 10% palladium on carbon for 16 hr. The catalyst is filtered off
25 through Celite filter aid, the filtrate concentrated, the residue taken up in ethyl acetate (300 ml)
and filtered again. The clear filtrate is concentrated to 200 ml and hexane added to a volume
of 450 ml. The cooled solution gives a small amount of gummy, dark material. The filtrate is
diluted with hexane and cooled overnight to give a solid. Recrystallization from ethyl
acetate/hexane then ethyl acetate gives 4-hydroxyindole-2-carboxamide, mp 219-221.

30 The 4-hydroxyindole-2-carboxamide (1.0 g), 1-chloro-3-[4-(diphenylmethyl)-1-p-
iperazinyl]propane (2.4 g) and powdered potassium carbonate (0.9 g) are combined in acetone
and heated at reflux overnight. The mixture is concentrated under reduced pressure, the
residue is treated with ice water and the whole extracted with methylene chloride (3 x 300 ml).
The pooled extract is washed with water, dried over magnesium sulfate and concentrated. The
35 residue is triturated with methylene chloride and recrystallized from methanol-ethyl acetate to
give the title compound, mp 222-3°; structure is supported by NMR, IR and mass spectra.

EXAMPLE 6 4-[3-[4-[2,4-Dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indole-2-carboxamide (VI)

4-(3-Chloropropoxy)indole-2-carboxamide (0.44 g), 4-(1-piperazinyl)-2,6-di-1-pyrrolidinylpyrimidine (0.525 g), powdered potassium carbonate (0.12 g) and sodium iodide (0.075 g) are combined in acetonitrile (50 ml) and heated at reflux for 96 hr. Although TLC showed both starting material and product remained in the reaction mixture, the mixture was concentrated under reduced pressure and the residue partitioned between 1N potassium bicarbonate and methylene chloride. The layers are separated, the aqueous layer is extracted with methylene chloride, the pooled organic extract is washed with water, then saline, dried over magnesium sulfate and concentrated. The residue is chromatographed over silica gel (250 ml) and eluted with methanol/methylene chloride (3.5/96.5). The appropriate fractions are pooled and concentrated. Crystallization from acetonitrile gives the title compound, mp 151-2° (foam); the proposed structure is supported by NMR, IR and mass spectra.

EXAMPLE 7 2-Cyano-4-[3-[4-(diphenylmethyl)-1-piperazinyl]propoxy]indole, dihydrochloride (VI)

4-[3-[4-(Diphenylmethyl)-1-piperazinyl] propoxy]indole-2-carboxamide (EXAMPLE 5, 1.45 g) and pyridine (1.18 ml) in dioxane (40 ml) is cooled to 12°. Trifluoroacetic anhydride (1.1 ml) in dioxane (8 ml) is added dropwise over 15 min (the reaction temperature remained between 12 and 14°). The mixture is stirred for 2 hr at 20-25°, diluted with methylene chloride and the mixture is washed with aqueous sodium bicarbonate, water and finally saline. The organic solution is dried over magnesium sulfate and concentrated to give a solid. The residue is dissolved in ether, filtered through magnesol, and the solution concentrated to give the free base of the title compound. The hydrochloride salt is prepared and recrystallized with an impurity. The solids and filtrates are combined, concentrated, and the residue partitioned between saturated sodium bicarbonate and methylene chloride to give the free base. This is chromatographed over silica gel (500 ml), eluting with methanol/methylene chloride (2.5/97.5). Twenty ml fractions are collected. Fractions 58-84 contained clean product. The hydrochloride salt is prepared from an ether solution by the addition of ethereal hydrochloric acid. Recrystallization from methanol/acetone/ether gives the title compound, mp 231-2°. The proposed structure is supported by NMR, IR and mass spectra.

EXAMPLE 8 2-Cyano-4-[3-[4-(2,4-dipyrrolidino-6-pyrimidinyl)-1-piperazinyl]propoxy]indole (VI)

The 4-hydroxyindole-2-carboxamide (EXAMPLE 5, 0.44 g), 4-(1-piperazinyl)-2,6-di-1-pyrrolidinylpyrimidine (0.525 g, 1.74 mmol), powdered potassium carbonate (0.12 g, 0.87 mmol) and sodium iodide (0.075 g) are combined in acetonitrile (50 ml) and heated at reflux for 96 hr. Although TLC showed starting material remained in the reaction mixture, the mixture is


concentrated under reduced pressure and the residue partitioned between 1N potassium bicarbonate and methylene chloride. The layers are separated, the aqueous layer is extracted with methylene chloride. The pooled organic extract is washed with water, then saline, dried over magnesium sulfate and concentrated. The residue is chromatographed over silica gel (250 ml) and eluted with methanol/methylene chloride (3.5/96.5). The appropriate fractions are pooled and concentrated; crystallization from acetonitrile gives 4-[3-[4-[2,4-dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indole-2-carboxamide, mp 151-2° (foam). The proposed structure is supported by NMR, IR and mass spectra.

A solution of 4-[3-[4-[2,4-dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indole-2-carboxamide (0.45 g) in dioxane (12 ml), under nitrogen, is mixed with pyridine (0.33 ml), cooled to 12° and treated during 5 min. with a solution of trifluoroacetic anhydride (0.31 ml) in dioxane (2.5 ml). The mixture is stirred at 12° for 15 min. and at 20-25° for 18 hr. It is again cooled to 12° and treated with additional trifluoroacetic anhydride (0.1 ml) in dioxane (1 ml). The mixture is kept at 12° for 15 min. and at 20-25° for 4 hr. It is then diluted with methylene chloride; washed with saturated sodium bicarbonate, water and saline, dried over magnesium sulfate and concentrated under reduced pressure. The residue is chromatographed over silica gel (275 ml) with eluting with ammonium hydroxide/methanol/methylene chloride (0.5/5/94.5). The product isolated from this column is crystallized from methanol to give the title compound, mp of an analytical sample 147-149°; Anal. calc'd for $C_{28}H_{36}N_8O$: C, 67.17; H, 7.25; N, 22.38. Found: C, 66.80; H, 7.34; N, 21.78.

-36-


CHART A
SENSITIZING AMINES

5




(I)

10



(II)

15




(III)

20

CHART A - Continued
SENSITIZING AMINES


5



(IV)


10

15



(V)

20




(VI)

-38-

CHART B


pyridin-2-,

5



(F-1)


10 3-,



(F-2)

15

or 4-yl optionally substituted
optionally as the N-oxide




(F-3)

20


$-*CH_2-(CH_2)_c-G-(CH_2)_d-CH_2-N^*-$

25



(F-4)

30 3-pyrrolin-1-yl



(F-5)

CHART B - Continued









- pyrrol-1-yl optionally substituted
- 5  $(C_1-C_3 \text{ alkyl})_{0-1}$ (F-6)
- piperidin-1-yl optionally substituted
- 10  $(C_1-C_3 \text{ alkyl})_{0-1}$
 $(C_1-C_3 \text{ alkyl})_{0-1}$ (F-7)
- 1,2,3,6-tetrahydropyridin-1-yl
- 15  (F-8)
- 1-hexamethyleneimino containing a 3- or 4- double bond or
- 20  (F-9)
- 1,4-dihydro-1-pyridinyl
- 25  a number of different groups
a number of different groups (F10)
- 1,3,5-triazin-2-yl or the
- 30  $(R_{21-2})_{0-1}$
 $(O)_{0-1}$
 $(R_{21-2})_{0-1}$ (F-11)
- N_1 -oxide thereof optionally substituted at the 4- and/or 6- position

CHART B - Continued


- 5 pyrimidin-4-yl or the N₁-oxide thereof optionally substituted at the 2- and/or 6- and 5- and/or 6- position




- 10 pyrimidin-2-yl optionally substituted




- 15 pyrazin-2-yl optionally substituted



- 20 imidazol-2-yl optionally substituted



- 25 1,3,4-triazol-2-yl optionally substituted



- 30 imidazol-4- or 5-yl optionally substituted




CHART B - Continued







- 5 benzo[b]thien-2-yl  (F-18)
- 10 indol-2-yl  (F-19)
- benzo[b]thiazol-2-yl  (F-20)
- 15 benzimidazol-2-yl  (F-21)
- 20 4-[2-[4-[2,6-di-(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl]  (F-22)
- 25 1,2,4-triazin-3-yl optionally substituted at the 5- and/or 6-position  (F-23)
- 30

CHART B - Continued




- (1-piperazinyl)-(C₂-C₄)
5 optionally substituted —(C₂-C₄ alkyl)—N—N—(Aryl or Heteroaryl) in the 4-position (F-24)
- 10 (1-piperazinyl)acetyl substituted in the 4-position —CO-CH₂—N—N—(Heteroaryl) (F-25)
- 15 (1-piperazinyl)carbonylmethyl substituted in the 4-position —CH₂-CO—N—N—(Heteroaryl) (F-26)
- 20 2-(carboxy)-1-pyrrolidiny  (F-27)
- 25 2-(carboxy)-1-piperidiny  (F-28)
- 30 2-(carboxy)-1-hexamethyleneimino  (F-29)

CHART B - Continued








- 5 2-(carboxy)-1-heptamethyleneimino  (F-30)
- 10 1-piperazinyl substituted in the 4-position  (F-31)
- 15 1-piperazinyl substituted in the 4-position  (F-32)
- 20 1-piperazinyl substituted in the 4-position  (F-33)
- 25 4-hydroxy-1-piperidiny1 substituted in the 4-position  (F-34)
- 30 1-piperazinyl substituted in the 4-position  (F-35)

CHART B - Continued


5 1-piperazinyl substituted
in the 4-position



(F-36)

10


15 1-piperazinyl
substituted in
the 4-position



(F-37)

CLAIMS

1. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing steroidal amine of formula (I)



(I)

where:

- 15 (A-I) R_6 is $\alpha\text{-}R_{6-1}:\beta\text{-}R_{6-2}$, R_{10} is $\alpha\text{-}R_{10-1}:\beta\text{-}R_{10-2}$ and R_7 is $\alpha\text{-}H:\beta\text{-}H$, where one of R_{6-1} and R_{6-2} is $-H$, and the other is $-H$, $-F$, or $C_1\text{-}C_3$ alkyl, R_{10-2} is $-\text{CH}_3$, R_{10-1} and R_5 taken together are $-(\text{CH}_2)_2\text{-C}(=\text{R}_{3-3})\text{-CH=}$ or $-\text{CH=CH-CO-CH=}$, where R_{3-3} is $=O$ or $\alpha\text{-}H:\beta\text{-OR}_{3-4}$ or $\alpha\text{-OR}_{3-4}:\beta\text{-H}$, where R_{3-4} is $-H$, $-\text{CO-CH}_3$, $-\text{CO-C}_2\text{H}_5$, $-\text{CO-C}_6\text{H}_5$, $-\text{CO-O-CH}_3$ or $-\text{CO-O-C}_2\text{H}_5$;
- 20 (A-II) R_5 is $\alpha\text{-}R_{5-3}:\beta\text{-}R_{5-4}$, R_6 is $\alpha\text{-}R_{6-3}:\beta\text{-}R_{6-4}$, R_{10} is $\alpha\text{-}R_{10-3}:\beta\text{-}R_{10-4}$ and R_7 is $\alpha\text{-}H:\beta\text{-}H$, where one of R_{6-3} and R_{6-4} is $-H$, and the other taken together with one of R_{5-3} and R_{5-4} forms a second bond between C_5 and C_6 , R_{10-4} is $-\text{CH}_3$, R_{10-3} and the other of R_{5-3} and R_{5-4} taken together is $-(\text{CH}_2)_2\text{-C(H)(OH)-CH}_2\text{-}$;
- (A-III) R_{10} and R_5 taken together are $=\text{CH-CH=C(OR}_3\text{)-CH=}$ where R_3 is
- 25 $-H$, $C_1\text{-}C_3$ alkyl, $-\text{CO-H}$, $C_2\text{-}C_4$ alkanoyl or benzyl, R_6 is $\alpha\text{-}R_{6-5}:\beta\text{-}R_{6-6}$ where one of R_{6-5} and R_{6-6} is $-H$, and the other is $-H$, $-F$, or $C_1\text{-}C_3$ alkyl and R_7 is $\alpha\text{-}H:\beta\text{-}H$;
- (A-IV) R_5 is $\alpha\text{-}R_{5-7}:\beta\text{-}R_{5-8}$, R_6 is $\alpha\text{-}R_{6-7}:\beta\text{-}R_{6-8}$, R_7 is $\alpha\text{-}H:\beta\text{-}H$ and R_{10} is $\alpha\text{-}R_{10-7}:\beta\text{-}R_{10-8}$, where one of R_{5-7} and R_{5-8} is $-H$, R_{10-7} and the other of R_{5-7} and R_{5-8} taken together are $-(\text{CH}_2)_2\text{-C}(=\text{R}_{3-3})\text{-CH}_2$, where R_{3-3} is as defined above, R_{10-8} is
- 30 $-\text{CH}_3$, where one of R_{6-7} and R_{6-8} is $-H$ and the other is $-H$, $-F$, or $C_1\text{-}C_3$ alkyl;
- (A-V) R_6 is $R_{6-9}:R_{6-10}$, R_7 is $R_{7-9}:R_{7-10}$, R_{10} is $\alpha\text{-}R_{10-9}:R_{10-10}$ where one of R_{6-9} and R_{6-10} is $-H$ and the other taken together with one of R_{7-9} and R_{7-10} forms a second bond between C_6 and C_7 , and the other of R_{7-9} and R_{7-10} is $-H$, R_{10-10} is $-\text{CH}_3$, R_{10-9} and R_5 taken together are $-(\text{CH}_2)_2\text{-C}(=\text{R}_{3-3})\text{-CH=}$ or $-\text{CH=CH-CO-CH=}$, where R_{3-3} is as defined
- 35 above;
- where:

(C-I) R_{11} is α - R_{11-1} : β - R_{11-2} , where one of R_{11-1} and R_{11-2} is taken together with R_9 to form a second bond between C_9 and C_{11} and the other of R_{11-1} and R_{11-2} is -H;

(C-II) R_9 is -Cl and R_{11} is =O or α -H: β - R_{11-4} where R_{11-4} is -Cl or -OH;

(C-III) R_9 is -H or -F and R_{11} is =O or α - R_{11-5} : β - R_{11-6} , where one of R_{11-5} and R_{11-6} is -H, and the other of R_{11-5} and R_{11-6} is -H, -OH or C_1 - C_{12} alkoxy;

(C-IV) R_9 is -H or -F and R_{11} is α -O-CO- R_{11-7} : β -H, where R_{11-7} is

(A) C_1 - C_3 alkyl,

(B) C_1 - C_{12} alkoxy,

(C) furanyl,

(D) - $NR_{122}R_{123}$, where one of R_{122} and R_{123} is -H, methyl or ethyl and the other is -H, C_1 - C_4 alkyl or phenyl,

(E) - X_3 -Aryl, where X_3 is -O- or a valence bond, where Aryl is phenyl optionally substituted with 1 thru 2 -Cl, -Br, C_1 - C_3 alkoxy, -COOH, - NH_2 , C_1 - C_3 alkylamino, di(C_1 - C_3)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2 - C_4 acylamino and -NH-CHO or with 1 -F or - CF_3 ;

where:

(D-I) R_{16} is R_{16-1} : R_{16-2} and R_{17} is R_{17-1} : R_{17-2} , where one of R_{16-1} and R_{16-2} is -H or - CH_3 and the other taken together with one of R_{17-1} and R_{17-2} forms a second bond between C_{16} and C_{17} , and the other of R_{17-1} and R_{17-2} is -C(=Z)-(CH₂)_n- $NR_{21-\alpha}R_{21-\beta}$ where Z is =O, =CH₂ or R_{17-9} :H where R_{17-9} is -H or - CH_3 , where n is 0 thru 6, where

(A) $R_{21-\alpha}$ is

(I) -(CH₂)_m- NR_{21-1} -Heteroaryl, where m is 2, 3 or 4, where R_{21-1} is -H or C_1 - C_3 alkyl, where Heteroaryl is:

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide

thereof optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is

(i) -F,

(ii) -Cl,

(iii) -Br,

(iv) C_1 - C_5 alkyl,


(v) -CH₂-CH=CH₂,

(vi) -Aryl, where Aryl is as defined above,

(vii) - $NR_{21-3}R_{21-3}$ where the R_{21-3} 's are the same or different and are -H, C_1 - C_3 alkyl or -CH₂-CH=CH₂,

(viii) *CH₂-(CH₂)_q-CH₂-N* - where the atoms marked

with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,



where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4), where G is -O-, -S-, -SO-, -SO₂- or -NHR₂₁₋₄, where R₂₁₋₄ is -H, C₁-C₃ alkyl, or Aryl as defined above, where c and d are the same or different and are 0 thru 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,

(ix) 3-pyrrolin-1-yl, (F-5)

(x) pyrrol-1-yl optionally substituted with C₁-C₃ alkyl,

10 (F-6)

(xi) piperidin-1-yl optionally substituted with 1 or 2 C₁-C₃ alkyl, (F-7)

(xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)

(xiii) 1-hexamethyleneimino containing a 3- or 4- double bond or 3- and 5- double bonds, (F-9)

(xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by two C₁-C₃ alkyl being the same or different, (F-10)

(xv) -OH,

(xvi) C₁-C₃ alkoxy,

20 (xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where R₂₁₋₇ is -H or C₁-C₃ alkyl and e is 0 thru 3,

(xviii) pyridin-2-, 3- or 4-yl,

(b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the 4- and/or 6- position with R₂₁₋₂ is as defined above, (F-11)

25 (c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2- and/or 6- position with R₂₁₋₂ is as defined above, (F-12)

(d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1 or 2 R₂₁₋₂ as is defined above, (F-13)

(e) pyrazin-2-yl optionally substituted with 1 or 2 R₂₁₋₂ as is defined above, (F-14)

(f) imidazol-2-yl optionally substituted in the 1 position with C₁-C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R₂₁₋₂ as defined above, (F-15)

35 (g) 1,2,4-triazol-3-yl optionally substituted in the 1 position with C₁-C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R₂₁₋₂ as defined above, (F-16)

(h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)

(i) benzo[b]thien-2-yl, (F-18)

5 (j) indol-2-yl, (F-19)

(k) benzo[b]thiazol-2-yl, (F-20)

(l) benzimidazol-2-yl, (F-21)

(m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl]piperazinyl, (F-22)

10 (n) 1,2,4-triazol-3-yl optionally substituted at the 5- and/or 6-position with R_{21-2} as is defined above, (F-23)

(2) (1-piperazinyl)-(C_2-C_4)alkyl optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)

(4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is

15 (a) $-O-CH_2CH_2-Y$, where Y is C_1-C_3 alkylamino, di(C_1-C_3)alkylamino where the alkyl groups are the same or different, C_3-C_6 alkyleneimino, optionally substituted with 1 or 2 C_1-C_3 alkyl,

(b) $-NR_{21-20}CH_2CH_2-Y$, where R_{21-20} is -H or C_1-C_3 alkyl and Y is as defined above,

20 (c) $-(CH_2)_g-N(R_{21-20})-Heteroaryl$, where g is 2, 3 or 4, and where R_{21-20} and Heteroaryl are as defined above,

(5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1-C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3-C_6 heterocyclic ring and
25 where m is as defined above,

(6) $-(CHCH_3)_b-(CH_2)_f-R_{21-24}$, where b is 0 and f is 1 thru 3 or b is one and f is 0 thru 3, where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1-C_3 alkoxy, - $NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1-C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4-C_7 cyclicamino ring,

30 (7) $-(CH_2)_i-Heteroaryl$, where i is 1 thru 4 and Heteroaryl is as defined above,

(8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-25)

(9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by
35 -Heteroaryl where Heteroaryl is as defined above, and (F-26)

(B) $R_{21-\beta}$ is

- (1) -H,
 (2) C₁-C₃ alkyl,
 (3) C₅-C₇ cycloalkyl,
 (4) -(CH₂)_m-NR₂₁₋₁-Heteroaryl, where m, R₂₁₋₁ and Heteroaryl are as
 5 defined above,
 (5) (1-piperazinyl)-(C₂-C₄)alkyl optionally substituted in the 4- position
 with -Aryl or -Heteroaryl as defined above, (F-24)
 (6) -(CH₂)_m-X₄, where m and X₄ are as defined above,
 (7) -(CH₂)_m-NR₂₁₋₂₂R₂₁₋₂₃, where m, R₂₁₋₂₂ and R₂₁₋₂₃ are as
 10 defined above,
 (8) -(CHCH₃)_b-(CH₂)_fR₂₁₋₂₄, where b, f and R₂₁₋₂₄ are as defined
 above,
 (C) R_{21-α} and R_{21-β} are taken together with the attached nitrogen atom to form
 a heterocyclic ring selected from the group consisting of
 15 (1) 2-(carboxy)-1-pyrrolidinyl optionally as the C₁-C₃ alkyl ester or as a
 pharmaceutically acceptable salt, (F-27)
 (2) 2-(carboxy)-1-piperidinyl optionally as the C₁-C₃ alkyl ester or as a
 pharmaceutically acceptable salt, (F-28)
 (3) 2-(carboxy)-1-hexamethyleneimino optionally as the C₁-C₃ alkyl
 20 ester or as a pharmaceutically acceptable salt, (F-29)
 (4) 2-(carboxy)-1-heptamethyleneimino optionally as the C₁-C₃ alkyl
 ester or as a pharmaceutically acceptable salt, (F-30)
 (5) 1-piperazinyl substituted in the 4- position with R₂₁₋₂₈-CO-(CH₂)_j-
 where R₂₁₋₂₈ is -Aryl, -NR₂₁₋₂₉Aryl and 2-furanyl, where R₂₁₋₂₉ is -H or C₁-C₃ alkyl, where
 25 j is 0 thru 3 and Aryl is as defined above, (F-31)
 (6) 1-piperazinyl substituted in the 4- position with Heteroaryl-(CH₂)_j-,
 where Heteroaryl and j are as defined above, (F-32)
 (7) 1-piperazinyl substituted in the 4- position with
 Aryl-(CH₂)_j-, where Aryl and j are as defined above, (F-33)
 30 (8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as
 defined above, (F-34)
 (9) 1-piperazinyl substituted in the 4- position with Heteroaryl-NR₂₁₋₂₉-
 CO-(CH₂)_i-, where Heteroaryl, R₂₁₋₂₉ and i are as defined above; (F-35)
 (D-II) R₁₆ is α-R₁₆₋₃:β-R₁₆₋₄ where one of R₁₆₋₃ and R₁₆₋₄ is -H and the other is -
 35 H, -F, -CH₃ or -OH, and R₁₇ is =CH-(CH₂)_p-NR_{21-α}R_{21-β}, where p is 1 or 2, where R_{21-α}
 and R_{21-β} are as defined above;

(D-III) R_{16} is α - R_{16-5} : β - R_{16-6} and R_{17} is α - R_{17-5} : β - R_{17-6} , where R_{16-5} is -H, -OH, -F or -CH₃ and R_{16-6} is -H, -OH, -F, or -CH₃, with the proviso that at least one of R_{16-5} and R_{16-6} is -H, where R_{17-5} is -H, -OH, -CH₃, -CH₂CH₃, C₂-C₇ alkanoyloxy or -O-CO-Aryl, where Aryl is as defined above, and where R_{17-6} is

- 5 -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , where Z, n, R_{21- α} and R_{21- β} are as defined above;

(D-IV) the 16,17-acetonide of a compound where R_{16-5} is -OH, R_{16-6} is -H, R_{17-5} is -OH and R_{17-6} is -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , where Z, n, R_{21- α} and R_{21- β} are as defined above;

with the following overall provisos that:

- 10 (I) one of R_{16-1} or R_{16-2} is taken together with one of R_{17-1} or R_{17-2} to form a second bond between C₁₆ and C₁₇, only when R_{10} is α - R_{10-1} : β - R_{10-2} , α - R_{10-3} : β - R_{10-4} , α - R_{10-7} : β - R_{10-8} or α - R_{10-9} : β - R_{10-10} ,
 (II) R_{17} is =CH-(CH₂)_p-NR_{21- α} R_{21- β} , only when R_{10} is α - R_{10-1} :
 β - R_{10-2} , α - R_{10-3} : β - R_{10-4} , α - R_{10-7} : β - R_{10-8} or α - R_{10-9} : β - R_{10-10} .
 15 (III) R_5 and R_{10} taken together are =CH-CH=C(OR₃)-CH=, only when R_{17} is α - R_{17-5} : β - R_{17-6} or the 16,17-acetonide of a compound where R_{16} is α -OH: β -H and R_{17} is α -OH: β -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , and
 (IV) R_5 is α - R_{5-7} : β - R_{5-8} , only when R_{17} is α - R_{17-5} : β - R_{17-6} or α -OH: β -C(=Z)-(CH₂)_n-NR_{21- α} R_{21- β} , or the 16,17-acetonide thereof; and pharmaceutically
 20 acceptable salts thereof.

2. A method according to claim 1 where treating is preventing resistance to the chemotherapeutic agents.

- 25 3. A method according to claim 1 where treating is overcoming resistance to the chemotherapeutic agents which already exists.

4. A method according to claim 1 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder
 30 carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.

5. A method according to claim 1 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol,
 35 colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

[S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,

(S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[[phenylamino]carbonyl]-opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-

5 benzofurancarboxamide,

(7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.

10 6. A method according to claim 1 where the steroidal amine (I) is selected from the group consisting of

17 α -hydroxy-21-[4-(2-pyridinyl)-1-piperazinyl]-pregna-4,9(11)-diene-3,20-dione,
21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

15 21-[4-(2,6-di-1-pyrrolidinyl)-4-pyrimidinyl-1-piperazinyl]pregna-1,4,9(11)-triene-3,20-dione,

21-[4-3,6-bis(diethylamino)-2-pyridinyl]-1-piperazinyl-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

20 21-[4-(4,6-di-1-pyrrolidinyl-1,3,5-triazin-2-yl)-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,


21-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

21-[4-(4,6-di-1-pyrrolidinyl-2-pyrimidinyl)-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione.

25

7. A method according to claim 1 where the effective amount of the steroidal amine (I) is from about 1 mg/kg/day to about 500 mg/kg/day.

8. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients
30 which comprises administering to that human an effective amount of a sensitizing alkyl amine of formula (II)



where:

n_2 is 3-14;

35 X_2 is -H,
-OH,

-O-CO-(C₁-C₄ alkyl),

-O-CO-H,

-O-CO-O-(C₁-C₄ alkyl),

(C₁-C₄) alkoxycarbonyl,

5 -O-CO-Aryl where Aryl is ϕ optionally substituted with 1 thru 3 of the following which may be the same or different:

-OH,

-OCH₃,

-F, -Cl, -Br, -CF₃,

10 -C₁-C₃ alkyl, and

-CO-R₅ where R₅ is

-OH,

-NH₂,

-NHR₆ where R₆ is

15 ϕ ,

C₁-C₃ alkyl and

-N(R₁₄)(R₁₅) where R₁₄ and R₁₅ are the same or

different and are C₁-C₃ alkyl,

-O-Aryl, where Aryl is as defined above,

20 -CH(OH)Aryl, where Aryl is as defined above,

Aryl, where aryl is as defined above;

(A) R_{21- α} is

(1) -(CH₂)_m-NR₂₁₋₁-Heteroaryl, where m is 2, 3 or 4, where R₂₁₋₁ is -H or C₁-C₃ alkyl, where Heteroaryl is:

25 (a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R₂₁₋₂, being the same or different, where R₂₁₋₂ is

(i) -F,

(ii) -Cl,

(iii) -Br,

30 (iv) C₁-C₅ alkyl,

(v) -CH₂-CH=CH₂,

(vi) -Aryl, where Aryl is phenyl optionally substituted with 1

through 2 -F, -Cl, -Br, C₁-C₃ alkoxy, -COOH, -NH₂, C₁-C₃ alkylamino, di(C₁-

C₃)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl,

35 1-hexamethylenimino, 1-heptamethylenimino, C₂-C₄ acylamino and -NH-CHO or with 1 -F or -CF₃;

(vii) $-\text{NR}_{21-3}\text{R}_{21-3}$ where the R_{21-3} s are the same or different and are $-\text{H}$, $\text{C}_1\text{-C}_3$ alkyl or $-\text{CH}_2\text{-CH=CH}_2$,

(viii α) $^*\text{CH}_2\text{-(CH}_2)_q\text{-CH}_2\text{-N}^*$ where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 through

5 5,

(viii β) $^*\text{CH}_2\text{-(CH}_2)_c\text{-G-(CH}_2)_d\text{-CH}_2\text{-N}^*$ where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4), where G is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$ or $-\text{NR}_{21-4}-$, where R_{21-4} is $-\text{H}$, $\text{C}_1\text{-C}_3$ alkyl, or Aryl as defined above, where c and d are the same or different and are 0 through 2 with the proviso
10 that the total number of ring carbon atoms is 4, 5 or 6,

(ix) 3-pyrrolin-1-yl, (F-5)

(x) pyrrol-1-yl optionally substituted with $\text{C}_1\text{-C}_3$ alkyl, (F-6)

(xi) piperidin-1-yl optionally substituted with 1 or 2 $\text{C}_1\text{-C}_3$ alkyl, (F-7)

(xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)

(xiii) 1-hexamethyleneimino containing a 3- or 4- double bond or 3- and 5- double bonds, (F-9)

(xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by two $\text{C}_1\text{-C}_3$ alkyl being the same or different, (F-10)

(xv) $-\text{OH}$,

(xvi) $\text{C}_1\text{-C}_3$ alkoxy,

(xvii) $-\text{NR}_{21-7}(\text{CH}_2)_e\text{-Q}$ where Q is 2-pyridinyl where R_{21-7} is $-\text{H}$ or $\text{C}_1\text{-C}_3$ alkyl and e is 0 through 3,

25 (xviii) pyridin-3- or 4-yl,

(xix) $-\text{CF}_3$,

(xx) $-\text{CCl}_3$,

(xxi) $-\text{SCH}_3$,

(b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the 4- and/or 6- position with R_{21-2} is as defined above, (F-11)

(c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2- and/or 6-, and 5- and/or 6- position with R_{21-2} is as defined above, (F-12)

(d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1 or 2 R_{21-2} as is defined above, (F-13)

35 (e) pyrazin-2-yl optionally substituted with 1 or 2 R_{21-2} as is defined above, (F-14)

- (f) imidazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-15)
- (g) 1,3,4-triazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)
- (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl], (F-22)
- (n) 1,2,4-triazin-3-yl optionally substituted at the 5- and/or 6- position with R_{M-2} as is defined above, (F-23)
- (2) $-(CH_2)_{2-4}-(1\text{-piperazinyl})$ optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)
- (3) -Heteroaryl, as defined above,
- (4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is
- (a) $-O-CH_2CH_2-Y$, where Y is C_1-C_3 alkylamino, di(C_1-C_3)alkylamino where the alkyl groups are the same or different, C_3-C_6 alkyleneimino, optionally substituted with 1 or 2 C_1-C_3 alkyl,
- (b) $-NR_{21-5}CH_2CH_2-Y$, where R_{21-5} is -H or C_1-C_3 alkyl and Y is as defined above,
- (c) $-(CH_2)_g-N(R_{21-5})\text{-Heteroaryl}$, where g is 2, 3 or 4, and where R_{21-5} and Heteroaryl are as defined above,
- (5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1-C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3-C_6 heterocyclic ring and where m is as defined above,
- (6) $-(CHCH_3)_b-(CH_2)_f\text{-Aryl}$ where b is 0 and f is 1 through 4 or b is 1 and f is 0 through 3, where Aryl is as defined above,
- (7) $-(CH_2)_i\text{-Heteroaryl}$, where i is 1 through 4 and Heteroaryl is as defined above,

(8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-25)

(9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-26)

5 (B) $R_{21-\beta}$ is

(1) -H,

(2) C_1-C_3 alkyl,

(3) C_5-C_7 cycloalkyl,

(4) $-(CH_2)_m-NR_{21-1}$ -Heteroaryl, where m, R_{21-1} and Heteroaryl are as defined above, 10

(5) (1-piperazinyl)-(C_2-C_4)alkyl optionally substituted in the 4- position with Aryl or Heteroaryl as defined above, (F-24)

(6) $-(CH_2)_m-X_4$, where m and X_4 are as defined above,

(7) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where m, R_{21-22} and R_{21-23} are as defined above, 15

(8) $-(CHCH_3)_b-(CH_2)_f-R_{21-24}$, where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1-C_3 alkoxy, $-NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1-C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4-C_7 cyclicamino ring and where b and f are as defined above,

20 (9) 2-pyridinylmethyl,

(C) $R_{21-\alpha}$ and $R_{21-\beta}$ are taken together with the attached nitrogen atom to form a heterocyclic ring selected from the group consisting of

(1) 2-(carboxy)-1-pyrrolidinyl optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-27)

25 (2) 2-(carboxy)-1-piperidinyl optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-28)

(3) 2-(carboxy)-1-hexamethyleneimino optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-29)

(4) 2-(carboxy)-1-heptamethyleneimino optionally as the C_1-C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-30)

(5) 1-piperazinyl optionally substituted in the 4- position with $R_{21-28}CO-(CH_2)_j$ - where R_{21-28} is -Aryl, -Heteroaryl, $-NR_{21-29}$ Heteroaryl and 2-furanyl, where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 through 3, and Aryl and Heteroaryl are as defined above, (F-31)

35 (6) 1-piperazinyl substituted in the 4- position with Heteroaryl- $(CH_2)_j$ -, where Heteroaryl and j are as defined above, (F-32)


- (7) 1-piperazinyl substituted in the 4- position with
Aryl-(CH₂)_j, where Aryl and j are as defined above, (F-33)
- (8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as defined
above, (F-34)
- 5 (9) 1-piperazinyl substituted in the 4- position with Heteroaryl-
NR₂₁₋₂₉-CO-(CH₂)_i, where Heteroaryl, R₂₁₋₂₉ and i are as defined above; (F-35)
- (10) 1-piperazinyl substituted in the 4- position with
-(CH₂)_j-C* = C(2-pyridinyl)-N=N-C(2-pyridinyl)=C*H, where * and j are as defined above, (F-36)
- 10 (11) 1-piperazinyl substituted in the 4- position with
-(CH₂)_i-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazine] (F-37)
and pharmaceutically acceptable salts thereof.
9. A method according to claim 8 where treating is preventing resistance to the
15 chemotherapeutic agents.
10. A method according to claim 8 where treating is overcoming resistance to the
chemotherapeutic agents which already exists.
- 20 11. A method according to claim 8 where the human cancer cells are selected from the group
consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder
carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple
myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.
- 25 12. A method according to claim 8 where the chemotherapeutic agents is selected from the
group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol,
colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine,
melphalan, adozelesin,
[S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-
30 3,6,7,8-tetrahydro-1-methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,
(S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[[[(phenylamino)carbonyl]-
opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-
benzofurancarboxamide,
(7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-
35 oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl]carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-
yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.

13. A method according to claim 8 where the alkyl amine (II) is selected from the group consisting of

- 4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinehexanol,
 4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazineoctanol,
 4-[[6-[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]hexyl]oxy]phenol,
 4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)- α -phenyl-1-piperazinebutanol,
 4-[3-[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]propyl]-2,5-dimethylphenol,
 4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazineheptanoic acid methyl ester.

14. A method according to claim 8 where the effective amount of the alkyl amine (II) is from about 1 mg/kg/day to about 500 mg/kg/day.

15. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing bicyclic amine of formula (III)



(III)

where:

- 25 W_2 is -O-, -S-, -NR₅₄- where R₅₄ is -H or C₁-C₃ alkyl,
 n_6 is 0, 1 or 2,
 R₇ is -H, C₁-C₄ alkyl, -CO-(C₁-C₄ alkyl), -CO- ϕ or -prodrug where prodrug is
 PO₂-O⁻ cation⁺ where cation⁺ is sodium, potassium or trialkylammonium where alkyl is C₁-
 C₃,
 30 -CO-CH₂-CO-NH-CH₂-SO₂-O⁻ cation⁺ where cation⁺ is as defined above,
 -CO-(CH₂)_{n₂₁}-R₅₁ where n₂₁ is 1-7 and R₅₁ is -COO⁻, -NR₅₁₋₁R₅₁₋₂ where
 R₅₁₋₁ and R₅₁₋₂ are the same or different and are -H or C₁-C₃ alkyl,
 -N⁺R₅₁₋₁R₅₁₋₂R₅₁₋₃ halide⁻ where R₅₁₋₁R₅₁₋₂R₅₁₋₃ are the same or different and are -H or
 C₁-C₃ alkyl, and where halide is -Cl or -Br,
 35 -CO-CH=CH-CO-O⁻ cation⁺ where cation⁺ is as defined above,
 -CO-N^{*}-CH=CH-N=C^{*}H where the atoms marked with an asterisk (*) are

bonded to each other resulting in the formation of a ring,

$-\text{CO}-\text{C}^*=\text{C}[(\text{CH}_2)_{n_{22}}-\text{NH}_2]-\text{CH}=\text{CH}-\text{CH}=\text{C}^*\text{H}$ where n_{22} is 1 or 2 and where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring,

5 $-\text{CO}-\text{C}^*=\text{CH}-\text{CH}=\text{C}(\text{NR}_{52})-\text{CH}=\text{C}^*\text{H}$ where R_{52} is -H or $\text{C}_1\text{-C}_3$ alkyl and where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring,

$-\text{CO}-(\text{CH}_2)_{n_{21}}-\text{CO}-\text{O}-[\text{C}_6\text{H}_{12}\text{O}_6 \text{ sugars}],$

$-\text{CO}-\text{O}-\text{CH}(\text{CH}_2-\text{O}-\text{CO}-\text{R}_{53})_2$ where the R_{53} 's are the same or different and are

10 $\text{C}_1\text{-C}_{18},$

$-\text{CO}-(\text{CH}_2)_6-\text{CO}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{SO}_3^- \text{ cation}^+$ where cation^+ is as defined

above,

$-\text{CH}_2-\text{O}-\text{CO}-(\text{CH}_2)_{n_{21}}-\text{NR}_{51-1}\text{R}_{51-2}$ where n_{21} , R_{51-1} and R_{51-2} are as defined

above,

15 $-\text{CO}-\text{NH}-\text{C}_6\text{H}_4-\text{R}_{55}$ where R_{55} is -H or $\text{C}_1\text{-C}_3$ alkyl, $-\text{NO}_2$,

$-\text{NR}_{51-1}\text{R}_{51-2}$ where R_{51-1} and R_{51-2} are as defined above and

R_{10} is -H or $-\text{CH}_3$,

R_{11} is -H or $-\text{CH}_3$,

R_{12} is -H or $-\text{CH}_3$,

20 (18-1) R_{16} is $\alpha\text{-R}_{16-1}\text{:}\beta\text{-R}_{16-2}$ where one of R_{16-1} and R_{16-2} is -H, $-\text{CH}_3$,

$-\text{CH}_2\text{CH}_3$ or $-\phi$ and the other is $-\text{X}_3-\text{NR}_{21-\alpha}\text{R}_{21-\beta}$ where X_3 is $-\text{CO}-$, $-(\text{CH}_2)_{n_{16}}-\text{CO}-$ where n_{16} is 1 or 2, $-(\text{CH}_2)_{n_3}-$ where n_3 is 1-6, or $-\text{CO}-\text{O}-(\text{CH}_2)_{n_{15}}-$ where n_{15} is 2-6, R_{25} and R_{26} are -H:-H;

(A) $R_{21-\alpha}$ is

25 (1) $-(\text{CH}_2)_m-\text{NR}_{21-1}\text{-Heteroaryl}$, where m is 2, 3 or 4, where R_{21-1} is -H or $\text{C}_1\text{-C}_3$ alkyl, where Heteroaryl is:

(a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof

optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is

(i) -F,

30

(ii) -Cl,

(iii) -Br,

(iv) $\text{C}_1\text{-C}_5$ alkyl,

(v) $-\text{CH}_2-\text{CH}=\text{CH}_2$,

(vi) -Aryl, where Aryl is phenyl optionally substituted with 1

35 thru 2 -F, -Cl, -Br, $\text{C}_1\text{-C}_3$ alkoxy, $-\text{COOH}$, $-\text{NH}_2$, $\text{C}_1\text{-C}_3$ alkylamino, $\text{di}(\text{C}_1\text{-C}_3)\text{alkylamino}$, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexa-

methylenimino, 1-heptamethylenimino, C_2-C_4 acylamino, $-NH-CHO$, with 1 $-F$ or $-CF_3$ or with 3,4-methylenedioxy and 3,4-ethylenedioxy;

(vii) $-NR_{21-3}R_{21-3}$ where the R_{21-3} 's are the same or different and are $-H$, C_1-C_3 alkyl or $-CH_2-CH=CH_2$,

5 (viii α) $^*CH_2-(CH_2)_q-CH_2-N^*$ - where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,

(viii β) $^*CH_2-(CH_2)_c-G-(CH_2)_d-CH_2-N^*$ - where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4), where G is $-O-$, $-S-$, $-SO-$, $-SO_2-$ or $-NR_{21-4}$, where R_{21-4} is $-H$, C_1-C_3 alkyl, or Aryl as
10 defined above, where c and d are the same or different and are 0 thru 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,

(ix) 3-pyrrolin-1-yl, (F-5)

(x) pyrrol-1-yl optionally substituted with

C_1-C_3 alkyl, (F-6)

15 (xi) piperidin-1-yl optionally substituted with 1 or 2

C_1-C_3 alkyl, (F-7)

(xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)

(xiii) 1-hexamethyleneimino containing a 3- or 4- double bond
or 3- and 5- double bonds, (F-9)

20 (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by
two C_1-C_3 alkyl being the same or different, (F-10)

(xv) $-OH$,

(xvi) C_1-C_3 alkoxy,

(xvii) $-NR_{21-7}(CH_2)_e-Q$ where Q is 2-pyridinyl where

25 R_{21-7} is $-H$ or C_1-C_3 alkyl and e is 0 thru 3,

(xviii) pyridin-2-, 3- or 4-yl,

(xix) $-CF_3$,

(xx) $-CCl_3$,

(xxi) $-SCH_3$,

30 (b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the
4- and/or 6- position with R_{21-2} is as defined above, (F-11)

(c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2-
and/or 6-, and 5- and/or 6- position with R_{21-2} is as defined
above, (F-12)

35 (d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1
or 2 R_{21-2} as is defined above, (F-13)

- (e) pyrazin-2-yl optionally substituted with 1 or 2 R_{21-2} as is defined above, (F-14)
- (f) imidazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-15)
- (g) 1,3,4-triazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)
- (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]ethyl], (F-22)
- (n) 1,2,4-triazin-3-yl optionally substituted at the 5- and/or 6- position with R_{21-2} as is defined above, (F-23)
- (2) $-(CH_2)_{2-4}$ -(1-piperazinyl) optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)
- (3) -Heteroaryl, as defined above,
- (4) $-(CH_2)_m-X_4$ where m is as defined above and where X_4 is
- (a) $-O-CH_2CH_2-Y$, where Y is C_1-C_3 alkylamino, di(C_1-C_3)alkylamino where the alkyl groups are the same or different, C_3-C_6 alkyleneimino, optionally substituted with 1 or 2 C_1-C_3 alkyl,
- (b) $-NR_{21-20}CH_2CH_2-Y$, where R_{21-20} is -H or C_1-C_3 alkyl and Y is as defined above,
- (c) $-(CH_2)_g-N(R_{21-20})$ -Heteroaryl, where g is 2, 3 or 4, and where R_{21-20} and Heteroaryl are as defined above,
- (5) $-(CH_2)_m-NR_{21-22}R_{21-23}$, where R_{21-22} is -H or C_1-C_3 alkyl and R_{21-23} is -Aryl or -Heteroaryl as defined above, or R_{21-22} and R_{21-23} are taken together with the attached nitrogen atom to form a saturated mono-nitrogen C_3-C_6 heterocyclic ring and where m is as defined above,
- (6) $-(CHCH_3)_b-(CH_2)_f$ -Aryl where b is 0 and f is 1 thru 4 or b is 1 and f is 0 thru 3, where Aryl is as defined above,

(7) $-(CH_2)_i$ -Heteroaryl, where i is 1 thru 4 and Heteroaryl is as defined above,

(8) (1-piperazinyl)acetyl substituted in the 4- position by Heteroaryl where Heteroaryl is as defined above, (F-25)

5 (9) (1-piperazinyl)carbonylmethyl substituted in the 4- position by -Heteroaryl where Heteroaryl is as defined above, and (F-26)

(B) $R_{21-\beta}$ is

(1) -H,

(2) C_1 - C_3 alkyl,

10 (3) C_5 - C_7 cycloalkyl,

(4) $-(CH_2)_m$ - NR_{21-1} -Heteroaryl, where m , R_{21-1} and Heteroaryl are as defined above,

(5) (1-piperazinyl)-(C_2 - C_4)alkyl optionally substituted in the 4- position with -Aryl or -Heteroaryl as defined above, (F-24)

15 (6) $-(CH_2)_m$ - X_4 , where m and X_4 are as defined above,

(7) $-(CH_2)_m$ - $NR_{21-22}R_{21-23}$, where m , R_{21-22} and R_{21-23} are as defined above,

(8) $-(CHCH_3)_b$ - $(CH_2)_f$ - R_{21-24} , where R_{21-24} is phenyl substituted with 1 thru 3 -OH, C_1 - C_3 alkoxy, $-NR_{21-25}R_{21-26}$ where R_{21-25} and R_{21-26} are the same or different and are -H, C_1 - C_3 alkyl or are taken together with the attached nitrogen atom to form a C_4 - C_7 cyclicamino ring and where b and f are as defined above,

(9) 2-pyridinylmethyl,

(10) 2-phenylethyl,

(C) $R_{21-\alpha}$ and $R_{21-\beta}$ are taken together with the attached nitrogen atom to form a heterocyclic ring selected from the group consisting of

(1) 2-(carboxy)-1-pyrrolidinyl optionally as the C_1 - C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-27)

(2) 2-(carboxy)-1-piperidinyl optionally as the C_1 - C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-28)

30 (3) 2-(carboxy)-1-hexamethyleneimino optionally as the C_1 - C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-29)

(4) 2-(carboxy)-1-heptamethyleneimino optionally as the C_1 - C_3 alkyl ester or as a pharmaceutically acceptable salt, (F-30)

(5) 1-piperazinyl optionally substituted in the 4- position with R_{21-28} -CO- $(CH_2)_j$ - where R_{21-28} is -Aryl, -Heteroaryl, $-NR_{21-29}$ -Heteroaryl and 2-furanyl, where R_{21-29} is -H or C_1 - C_3 alkyl, where j is 0 thru 3, and Aryl and Heteroaryl are as defined above, (F-31)

(6) 1-piperazinyl substituted in the 4-position with Heteroaryl(CH₂)_j- where Heteroaryl and j are as defined above (F-32)

(7) 1-piperazinyl substituted in the 4-position with Aryl-(CH₂)_j-, where Aryl and j are as defined above, (F-33)

5 (8) 4-hydroxy-1-piperidinyl substituted in the 4- position with Aryl as defined above, (F-34)

(9) 1-piperazinyl substituted in the 4- position with Heteroaryl-NR₂₁₋₂₉-CO-(CH₂)_i-, where Heteroaryl, R₂₁₋₂₉ and i are as defined above; (F-35)

10 (10) 1-piperazinyl substituted in the 4- position with -(CH₂)_j-C^{*}=C(2-pyridinyl)-N=N-C(2-pyridinyl)=C^{*}H, where * and j are as defined above, (F-36)

(11) 1-piperazinyl substituted in the 4- position with -(CH₂)_i-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazine] where i is as defined above, (F-37)

15 (12) 1-piperazinyl substituted in the 4- position with C₁-C₃ alkyl optionally substituted with 1 or 2 Aryl;

(18-2) n₆ is 0, R₁₆ is R₁₆₋₃:R₁₆₋₄ and R₂₅ is R₂₅₋₃:R₂₅₋₄ where one of R₁₆₋₃ and R₁₆₋₄ is taken together with one of R₂₅₋₃ and R₂₅₋₄ to form a second bond between the carbon atoms to which R₁₆ and R₂₅ are attached and the other of R₁₆₋₃ and R₁₆₋₄ is -X₃-NR_{21-α}R_{21-β} where X₃, R_{21-α} and R_{21-β} are as defined above and the other of R₂₅₋₃ and R₂₅₋₄ is -H, 20

(18-3) n₆ is 1, R₂₅ is R₂₅₋₅ and R₂₅₋₆ and R₂₆ is R₂₆₋₅ and R₂₆₋₆ where one of R₂₅₋₅ and R₂₅₋₆ and one of R₂₆ is R₂₆₋₅ and R₂₆₋₆ are taken together to form a second bond between the carbon atoms to which R₂₅ and R₂₆ are attached and the other of R₂₅₋₅ and R₂₅₋₆ and R₂₆₋₅ and R₂₆₋₆ are -H, and pharmaceutically acceptable salts thereof.

25

16. A method according to claim 15 where treating is preventing resistance to the chemotherapeutic agents.

17. A method according to claim 15 where treating is overcoming resistance to the chemotherapeutic agents which already exists. 30

18. A method according to claim 15 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma. 35

19. A method according to claim 15 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

5 [S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,

(S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[(phenylamino)carbonyl]-opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-benzofurancarboxamide,

10 (7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.

20. A method according to claim 15 where the bicyclic amine (I) is selected from the group
15 consisting of

2-[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]methyl]-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol,


2-[[4-3-(ethylamino)-2-pyridinyl]-1-piperazinyl]methyl]-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol,

20 3-[4-(2,6-di-1-pyrrolinyl-4-pyrimidinyl)-1-piperazinyl]-6-(acetyloxy)-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-2-carboxylic acid propyl ester,

1-[3-(ethylamino)-2-pyridinyl]-4-[(5-methoxy-4,6,7-trimethyl-1H-indol-2-yl)carbonyl]-piperazine.

25 21. A method according to claim 15 where the effective amount of the steroidal amine (I) is from about 1 mg/kg/day to about 500 mg/kg/day.

22. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing
30 bicyclic ether of formula (IV)



(IV)

5

where R_1 is -H or $-OR_{1-1}$ where R_{1-1} is C_1-C_3 alkyl and where R_2 is -H or $-OR_{2-1}$ where R_{2-1} is C_1-C_3 alkyl with the proviso that R_2 is not -H only when R_1 is -H,

10 R_3 is $\alpha-R_{3-1}:\beta-R_{3-2}$ where R_{3-1} and R_{3-2} are the same or different and are -H or $-CH_3$ with the proviso that R_{3-2} is not $-CH_3$ unless R_{3-1} is $-CH_3$,

n is 1, 2 or 3,

R_4 is $R_{4-1}:R_{4-2}$ where R_{4-1} is -H, $-CH_3$, $-CH_2CH_3$, 4-fluorophenyl, 4-chlorophenyl, R_{4-2} is $-(CH_2)_n-R_{4-3}$ where n is 1, 2 or 3 and where R_{4-3} is

15

-Cl,

1-piperazinyl optionally substituted in the 4-position with a member selected from the group consisting of

- ϕ optionally substituted with 1 - CF_3 , -Cl, -F, $-CH_3$, $-CH_2CH_3$,

2-pyridinyl optionally substituted in the 6-position with $-NR_{4-4}R_{4-5}$

20 where R_{4-4} and R_{4-5} are the same or different and are -H, C_1-C_3 alkyl and where R_{4-4} and R_{4-5} are taken together with the attached nitrogen atom to form a ring selected from the group consisting of 1-pyrrolidinyl, 1-piperidinyl, 1-morpholinyl,

4-pyrimidinyl optionally substituted in the 2 and/or 6-position with -

$NR_{4-4}R_{4-5}$ where R_{4-4} and R_{4-5} are as defined above,

25

piperid-3-en-1-yl optionally substituted in the 4-position with a

member selected from the group consisting of

- ϕ optionally substituted with 1 - CF_3 , -Cl, -F, $-CH_3$, $-CH_2CH_3$,

2-pyridinyl optionally substituted in the 6-position with $-NR_{4-4}R_{4-5}$

where R_{4-4} and R_{4-5} are the same or different and are -H, C_1-C_3 alkyl and where R_{4-4} and R_{4-5}

30 R_{4-5} are taken together with the attached nitrogen atom to form a ring selected from the group consisting of 1-pyrrolidinyl, 1-piperidinyl, 1-morpholinyl,

4-pyrimidinyl optionally substituted in the 2 and/or 6-position with -

$NR_{4-4}R_{4-5}$ where R_{4-4} and R_{4-5} are as defined above, and pharmaceutically acceptable salts thereof.

35

23. A method according to claim 22 where treating is preventing resistance to the chemotherapeutic agents.
24. A method according to claim 22 where treating is overcoming resistance to the
- 5 chemotherapeutic agents which already exists.
25. A method according to claim 22 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple
- 10 myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.
26. A method according to claim 22 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine,
- 15 melphalan, adozelesin,
- [S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,
- (S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[[phenylamino]carbonyl]-opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-
- 20 benzofurancarboxamide,
- (7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.
- 25 27. A method according to claim 22 where the bicyclic ether (IV) is selected from the group consisting of
- 1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-[3-(tribluoromethyl)-phenyl]piperazine,
- 1-[(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)-
- 30 piperazine,
- 1-[(3,4-dihydro-8-methoxy-1H-2-benzopyran-1-yl)methyl]-4-(2-methylphenyl)piperazine,
- 4-(4-chlorophenyl)-1-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1,2,3,6-tetrahydropiperidine,
- 1-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)-
- 35 piperazine,
- 1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-

- yl)ethyl]piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(4-fluorophenyl)piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-[3-(trifluoromethyl)phenyl]piperazine,
5 1-(4-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]piperazine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,
10 1-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,6-tetrahydro-4-phenylpyridine,
2-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,5-tetrahydro-6,7-dimethoxyisoquinoline,
1-[2-(3,4-dihydro-5,6-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]-4-phenylpiperazine,
15 1-(4-fluorophenyl)-4-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]-propyl]piperazine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-(2-methylphenyl)piperazine,
20 1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-phenylpiperazine,
2H-Benzimidazol-2-one, 1-[1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]-propyl]-4-piperidinyl]-1,3-dihydropiperazine,
1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl]propyl]-4-(2-pyridinyl)piperazine,
25 1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-1,2,3,6-tetrahydro-4-phenylpyridine,
1-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]-4-(2-methylphenyl)piperazine,
30 1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]piperazine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-phenylpiperazine,
1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-(2-methylphenyl)piperazine,
35 1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-4-(4-

fluorophenyl)piperazine,

1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl]propyl]-4-phenylpiperazine,

1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,

1-[3-(3,4-dihydro-6,7-dimethoxy-1,4-dimethyl-1H-2-benzopyran-1-yl)propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,

1-[3-[1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl]propyl]-4-phenylpiperazine,

1-(3-chloropropyl)-1-(4-fluorophenyl)-3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran,

1-[3-(3,4-dihydro-6,7-dimethoxy-1,4,4-trimethyl-1H-2-benzopyran-1-yl)propyl]-1,2,3,6-tetrahydro-4-phenylpyridine,

1-[3-(3,4-dihydro-6,7-dimethoxy-1,4-dimethyl-1H-2-benzopyran-1-yl)propyl]-4-(2-methylphenyl)piperazine,

1-(2-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-1-methyl-1H-2-benzopyran-1-yl)ethyl]piperazine,

1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4,4-dimethyl-1H-2-benzopyran-1-yl)ethyl]piperazine,

1-(3-chlorophenyl)-4-[2-(3,4-dihydro-6,7-dimethoxy-4-methyl-1H-2-benzopyran-1-yl)ethyl]piperazine,


6-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-N,N,N',N'-tetraethyl-2,4-pyrimidinediimine,

4-[4-[2-(3,4-dihydro-6,7-dimethoxy-1H-2-benzopyran-1-yl)ethyl]-1-piperazinyl]-2,6-di-1-pyrrolidinylpyrimidine.

28. A method according to claim 22 where the effective amount of the bicyclic ether (IV) is from about 1 mg/kg/day to about 500 mg/kg/day.

29. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing tricyclic compound of formula (V)

(V)



where:

n_1 is 1 thru 3,

R_1 is α - R_{1-1} : β - R_{1-2} where R_{1-1} and R_{1-2} are the same or different and are -H,

C_1 - C_3 alkyl,

R_2 is C_1 - C_3 alkyl,

R_3 is - ϕ optionally substituted with 1 thru 3 -F, -Cl, C_1 - C_3 alkyl and pharmaceutically acceptable salts thereof.

30. A method according to claim 29 where treating is preventing resistance to the chemotherapeutic agents.

31. A method according to claim 29 where treating is overcoming resistance to the chemotherapeutic agents which already exists.

32. A method according to claim 29 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.

33. A method according to claim 29 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

[S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b;4,3-b']dipyrrol-4-ol,

(S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[[[(phenylamino)carbonyl]-oxy]benzo[1,2-b;4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-

benzofurancarboxamide,

(7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-

oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl)carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.

34. A method according to claim 29 where the tricyclic compound (V) is selected from the group consisting of

1-(4-fluorophenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine,

1-(2-methylphenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine,

- 10 1-(2-chlorophenyl)-4-[2-(3a,4,5,6-tetrahydro-7-methoxy-4,4-dimethyl-1H-3H-naphtho[1,8-cd]pyran-1-yl)-ethyl]piperazine.

35. A method according to claim 24 where the effective amount of the tricyclic compound (V) is from about 1 mg/kg/day to about 500 mg/kg/day.

15

36. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a compound selected from the group consisting of

- 20 11-[[[4-(2,6-di-1-pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl]acetyl]oxy]-16 α -methylpregn-4-ene-3,20-dione,

4-[5-(benzoyloxy)-2,6-di-1-pyrrolidinyl-4-pyrimidinyl]1-piperazineheptanoic acid methyl ester,

21-[4-(2,6-di-1-pyrrolidinyl-5-(4-chlorobenzoyloxy))-4-pyrimidinyl-1-piperazinyl]-16 α -methylpregna-1,4,9(11)-triene-3,20-dione,

- 25 3-[2-[4-[3-(ethylamino)-2-pyridinyl]-1-piperazinyl]ethyl]octahydro-7-[2-(5-hydroxy-2-methylphenyl)ethyl]-3a-methyl-5H-inden-5-one and pharmaceutically acceptable salts thereof.

37. A method according to claim 36 where treating is preventing resistance to the chemotherapeutic agents.

30

38. A method according to claim 36 where treating is overcoming resistance to the chemotherapeutic agents which already exists.

39. A method according to claim 36 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple
- 35

myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.

40. A method according to claim 36 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

[S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-,methyl-benzo[1,2-b;4,3-b']dipyrrol-4-ol,

- (S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[(phenylamino)carbonyl]-opxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-benzofurancarboxamide,


(7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.

15

41. A method according to claim 36 where the effective amount of the compound is from about 1 mg/kg/day to about 500 mg/kg/day.

42. A method of treating resistance to cancer chemotherapeutic agents in human cancer patients which comprises administering to that human an effective amount of a sensitizing indole of formula (VI)

25



(VI)

30 where:

R_1 is $-C \equiv N$ or $-CO-NH_2$;

n_1 is 1 thru 5;

R_{21} is

(1) $R_{21-28}-CO-(CH_2)_j-$ where R_{21-28} is -Aryl, $-NR_{21-29}$ Aryl and 2-furanyl,

- 35 where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 thru 3 and Aryl is phenyl optionally substituted with 1 or 2 -Cl, -Br, C_1-C_3 alkoxy, $-COOH$, $-NH_2$, C_1-C_3 alkylamino, $di(C_1-$

C_3)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2 - C_4 acylamino and -NH-CHO or with 1 -F or -CF₃; (F-31)

(2) Heteroaryl-(CH₂)_j-, where Heteroaryl is

- 5 (a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R₂₁₋₂, being the same or different, where R₂₁₋₂ is
- (i) -F,
 - (ii) -Cl,
 - (iii) -Br,
 - 10 (iv) C₁-C₅ alkyl,
 - (v) -CH₂-CH=CH₂,
 - (vi) -Aryl, where Aryl is as defined above,
 - (vii) -NR₂₁₋₃R₂₁₋₃ where the R₂₁₋₃'s are the same or different and are -H, C₁-C₃ alkyl or -CH₂-CH=CH₂,
 - 15 (viiiα) *CH₂-(CH₂)_q-CH₂-N* - where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,
 - (viiiβ) *CH₂-CH₂-(CH₂)_c-G-(CH₂)_d-CH₂-CH₂-N* - where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring (F-4), where G is -O-, -S-, -SO₂- or -NHR₂₁₋₄, where R₂₁₋₄ is -H, C₁-C₃ alkyl, or Aryl
 - 20 as defined above, where c and d are the same or different and are 0 thru 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,
 - (ix) 3-pyrrolin-1-yl, (F-5)
 - (x) pyrrol-1-yl optionally substituted with C₁-C₃ alkyl, (F-6)
 - (xi) piperidin-1-yl optionally substituted with 1 or 2 C₁-C₃
 - 25 alkyl, (F-7)
 - (xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)
 - (xiii) 1-hexamethyleneimino containing a 3- or 4- double bond or 3- and 5- double bonds, (F-9)
 - (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by
 - 30 two C₁-C₃ alkyl being the same or different, (F-10)
 - (xv) -OH,
 - (xvi) C₁-C₃ alkoxy,
 - (xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where R₂₁₋₇ is -H or C₁-C₃ alkyl and e is 0 thru 3,
 - 35 (xviii) pyridin-2-, 3- or 4-yl,
- (b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the

- 4- and/or 6- position with R_{21-2} is as defined above, (F-11)
- (c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2- and/or 6- position with R_{21-2} is as defined above, (F-12)
- (d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1 or 2 R_{21-2} as is defined above, (F-13)
- (e) pyrazin-2-yl optionally substituted with 1 or 2 R_{21-2} as is defined above, (F-14)
- (f) imidazol-2-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-15)
- (g) 1,2,4-triazol-3-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R_{21-2} as defined above, (F-16)
- (h) imidazol-4- or 5-yl optionally substituted in the 1 position with C_1-C_3 alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R_{21-2} as defined above, (F-17)
- (i) benzo[b]thien-2-yl, (F-18)
- (j) indol-2-yl, (F-19)
- (k) benzo[b]thiazol-2-yl, (F-20)
- (l) benzimidazol-2-yl, (F-21)
- (m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]-ethyl]piperazinyl, (F-22)
- (n) 1,2,4-triazol-3-yl optionally substituted at the 5- and/or 6- position with R_{21-2} as is defined above, (F-23)
- and where j is as defined above, (F-32)
- (3) Aryl-(CH₂)_j-, where Aryl and j are as defined above, (F-33)
- and pharmaceutically acceptable salts thereof.

43. A method according to claim 42 where treating is preventing resistance to the chemotherapeutic agents.

44. A method according to claim 42 where treating is overcoming resistance to the chemotherapeutic agents which already exists.

45. A method according to claim 42 where the human cancer cells are selected from the group consisting of ovarian, sarcoma, non-Hodgkin's lymphoma, lung, breast cancer, bladder

carcinoma, colon carcinoma, pancreatic carcinoma, carcinoma of the ampulla of Vater, multiple myeloma, adult acute lymphocytic leukemia, adult non-lymphocytic leukemia and neuroblastoma.

46. A method according to claim 42 where the chemotherapeutic agents is selected from the group consisting of doxorubicin, daunomycin, vinca alkaloids, vincristine, vinblastine, taxol, colchicine, epipodophyllotoxins such as etoposide, actinomycin D, puromycin, emetine, melphalan, adozelesin,

- [S-(R,R)] 6,6'-[carbonylbis(imino-1H-indole-05,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-methyl-benzo[1,2-b:4,3-b']dipyrrol-4-ol,
- 10 (S)-N-[2-[[1-(chloromethyl)-1,6-dihydro-8-methyl-5-[(phenylamino)carbonyl]oxy]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1H-indol-5-yl]-6-(diethylamino)-2-benzofurancarboxamide,
- (7bR,8aS)-7-[[1,6-dihydro-4-hydroxy-5-methoxy-7-[(4,5,8,8a-tetrahydro-7-methyl-4-oxocyclopropa[c]pyrrolo[3,2-e]indol-2(1H)-yl)carbonyl]benzo[1,2-b:4,3-b']dipyrrol-3(2H)-yl]carbonyl]-1,6-dihydro-4-hydroxy-5-methoxybenzo[1,2-b:4,3-b']dipyrrole-3(2H)-carboxamide.
- 15


47. A method according to claim 42 where the indole (VI) is selected from the group consisting of

- 4-[3-4-diphenylmethyl)-1-piperazinyl]propoxy]indole-2-carboxamide,
- 20 4-[3-[4-[2,4-dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indol-2-carboxamide,
- 2-cyano-4-[3-[4-(diphenylmethyl)-1-piperazinyl]propoxy]indole,
- 2-cyano-4-[3-[4-(2,4-dipyrrolidino-6-pyrimidinyl)-1-piperazinyl]propoxy]indole.

48. A method according to claim 42 where the effective amount of the indole (VI) is from about 1 mg/kg/day to about 500 mg/kg/day.
- 25

49. An indole of formula (VI)

30



(VI)

35

where R_1 is $-C \equiv N$ or $-CO-NH_2$;

n_1 is 1 thru 5;

R_{21} is

- (1) $R_{21-28}-CO-(CH_2)_j-$ where R_{21-28} is -Aryl, $-NR_{21-29}$ Aryl and 2-furanyl,
 5 where R_{21-29} is -H or C_1-C_3 alkyl, where j is 0 thru 3 and Aryl is phenyl optionally substituted with 1 or 2 -Cl, -Br, C_1-C_3 alkoxy, -COOH, $-NH_2$, C_1-C_3 alkylamino, di(C_1-C_3)alkylamino, where the alkyl groups are the same or different, 1-pyrrolidinyl, 1-piperidinyl, 1-hexamethylenimino, 1-heptamethylenimino, C_2-C_4 acylamino and $-NH-CHO$ or with 1 -F or $-CF_3$; (F-31)
- 10 (2) Heteroaryl- $(CH_2)_j-$, where Heteroaryl is
 (a) pyridin-2- (F-1), 3- (F-2) or 4-yl (F-3) or the N-oxide thereof optionally substituted by 1 or 2 R_{21-2} , being the same or different, where R_{21-2} is
 (i) -F,
 (ii) -Cl,
 15 (iii) -Br,
 (iv) C_1-C_5 alkyl,
 (v) $-CH_2-CH=CH_2$,
 (vi) -Aryl, where Aryl is as defined above,
 (vii) $-NR_{21-3}R_{21-3}$ where the R_{21-3} 's are the same or different
 20 and are -H, C_1-C_3 alkyl or $-CH_2-CH=CH_2$,
 (viiiα) $^*CH_2-(CH_2)_q-CH_2-N^*$ where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring, where q is 1 thru 5,
 (viiiβ) $^*CH_2-CH_2-(CH_2)_c-G-(CH_2)_d-CH_2-CH_2-N^*$ where the atoms marked with an asterisk (*) are bonded to each other resulting in the formation of a ring
 25 (F-4), where G is -O-, -S-, -SO-, $-SO_2-$ or $-NHR_{21-4}$, where R_{21-4} is -H, C_1-C_3 alkyl, or Aryl as defined above, where c and d are the same or different and are 0 thru 2 with the proviso that the total number of ring carbon atoms is 4, 5 or 6,
 (ix) 3-pyrrolin-1-yl, (F-5)
 (x) pyrrol-1-yl optionally substituted with C_1-C_3 alkyl, (F-6)
 30 (xi) piperidin-1-yl optionally substituted with 1 or 2 C_1-C_3 alkyl, (F-7)
 (xii) 1,2,3,6-tetrahydropyridin-1-yl, (F-8)
 (xiii) 1-hexamethyleneimino containing a 3- or 4- double bond or 3- and 5- double bonds, (F-9)
 35 (xiv) 1,4-dihydro-1-pyridinyl substituted in the 4 position by two C_1-C_3 alkyl being the same or different, (F-10)

(xv) -OH,

(xvi) C₁-C₃ alkoxy,

(xvii) -NR₂₁₋₇-(CH₂)_e-Q where Q is 2-pyridinyl where R₂₁₋₇ is -H or C₁-C₃ alkyl and e is 0 thru 3,

5

(xviii) pyridin-2-, 3- or 4-yl,

(b) 1,3,5-triazin-2-yl or the N-oxide thereof optionally substituted at the 4- and/or 6- position with R₂₁₋₂ is as defined above, (F-11)

(c) pyrimidin-4-yl or the N-oxide thereof optionally substituted at the 2- and/or 6- position with R₂₁₋₂ is as defined above, (F-12)

10

(d) pyrimidin-2-yl optionally substituted at 4- and/or 6- position with 1 or 2 R₂₁₋₂ as is defined above, (F-13)

(e) pyrazin-2-yl optionally substituted with 1 or 2 R₂₁₋₂ as is defined above, (F-14)

15

(f) imidazol-2-yl optionally substituted in the 1 position with C₁-C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R₂₁₋₂ as defined above, (F-15)

(g) 1,2,4-triazol-3-yl optionally substituted in the 1 position with C₁-C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with R₂₁₋₂ as defined above, (F-16)

20

(h) imidazol-4- or 5-yl optionally substituted in the 1 position with C₁-C₃ alkyl or -Aryl, where Aryl is as defined above, and further optionally substituted with 1 or 2 R₂₁₋₂ as defined above, (F-17)

(i) benzo[b]thien-2-yl, (F-18)

(j) indol-2-yl, (F-19)

25

(k) benzo[b]thiazol-2-yl, (F-20)

(l) benzimidazol-2-yl, (F-21)

(m) 4-[2-[4-[2,6-bis(1-pyrrolidinyl)-4-pyrimidinyl]-1-piperazinyl]-ethyl]piperazinyl, (F-22)

30

(n) 1,2,4-triazol-3-yl optionally substituted at the 5- and/or 6- position with R₂₁₋₂ as is defined above, (F-23)

and where j is as defined above, (F-32)

(3) Aryl-(CH₂)_j, where Aryl and j are as defined above, (F-33)
and pharmaceutically acceptable salts thereof.

35 50. A indole according to claim 49 which is selected from the group consisting of
4-[3-4-diphenylmethyl)-1-piperazinyl]propoxy]indole-2-carboxamide,

4-[3-[4-[2,4-dipyrrolidino-6-pyrimidinyl]-1-piperazinyl]propoxy]indol-2-carboxamide,
2-cyano-4-[3-[4-(diphenylmethyl)-1-piperazinyl]propoxy]indole,
2-cyano-4-[3-[4-(2,4-dipyrrolidino-6-pyrimidinyl)-1-piperazinyl]propoxy]indole.